# abcam

## Product datasheet

# Anti-STUB1/CHIP antibody ab2917

★★★★★ 2 Abreviews 14 References 3 Images

Overview

Product name Anti-STUB1/CHIP antibody

**Description** Rabbit polyclonal to STUB1/CHIP

Host species Rabbit

Specificity Detects human carboxyl terminus of hsc70-interacting protein (CHIP). Detects a band of

approximately 35 kDa representing CHIP from COS-1 cells overexpressing the human gene including a non-specific band at 80kDa . A customer reported that in mouse cerebrum the antibody detects a band of 35kDa as well as two small mon-specific at 44 and 70 kDa.

Tested applications Suitable for: WB

Species reactivity Reacts with: Mouse, Human

Predicted to work with: Chicken

**Immunogen** Synthetic peptide corresponding to Human STUB1/CHIP aa 218-232.

Sequence:

**VDEKRKKRDIPDYLC** 

(Peptide available as ab4934)

Run BLAST with
Run BLAST with

Positive control WB: mouse brain, MCF-7 cells, transfected COS-1 cells

**General notes**The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As  $\,$ 

**Properties** 

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -

80°C. Avoid freeze / thaw cycle.

**Storage buffer** Preservative: 0.05% Sodium azide

1

Constituents: 0.1% BSA, 99% PBS

Purity Immunogen affinity purified

Primary antibody notes A recently identified protein, termed carboxyl terminus of hsc70-interacting protein (CHIP), has

been shown to interact both with the constitutive form of hsc70 and the stress inducible form, hsp70. This novel 35 kDa cytoplasmic protein has been shown to be highly expressed in striated muscle in vivo. Additional studies have shown that this protein is expressed over a broad range of cultured tissues. Through immunoprecipitation experiments, CHIP has been shown to directly bind to the carboxyl terminus of hsc70 and hsp70 where it decreases ATPase activity and reduces overall chaperone efficiency. CHIP has also been identified as an important protein in the

 $ubiquitin\hbox{-proteasome system. CHIP contains a $U$-box domain and acts as an E3 ubiquitin-ligase$ 

in conjunction with hsc70 and hsp90.

**Clonality** Polyclonal

**Isotype** IgG

#### **Applications**

The Abpromise guarantee Our Abpromise guarantee covers the use of ab2917 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB	<b>★★★★☆</b> (2)	Use at an assay dependent concentration. Predicted molecular weight: 35 kDa.

#### **Target**

**Function** E3 ubiquitin-protein ligase which targets misfolded chaperone substrates towards proteasomal

degradation. Ubiquitinates NOS1 in concert with Hsp70 and Hsp40. Modulates the activity of several chaperone complexes, including Hsp70, Hsc70 and Hsp90. Mediates transfer of non-canonical short ubiquitin chains to HSPA8 that have no effect on HSPA8 degradation. Mediates polyubiquitination of DNA polymerase beta (POLB) at 'Lys-41', 'Lys-61' and 'Lys-81', thereby playing a role in base-excision repair: catalyzes polyubiquitination by amplifying the HUWE1/ARF-BP1-dependent monoubiquitination and leading to POLB-degradation by the proteasome.

 $\label{lem:mediates} \textit{Mediates polyubiquitination of CYP3A4}.$ 

Tissue specificity Highly expressed in skeletal muscle, heart, pancreas, brain and placenta. Detected in kidney, liver

and lung.

Pathway Protein modification; protein ubiquitination.

**Sequence similarities** Contains 3 TPR repeats.

Contains 1 U-box domain.


**Domain** The TPR domain is essential for ubiquitination mediated by UBE2D1.

**Post-translational** Phosphorylated upon DNA damage, probably by ATM or ATR. **modifications** Auto-ubiquitinated; mediated by UBE2D1 and UBE2D2.


Auto-ubiquitifiated, mediated by OBEZD I and OBEZDZ:

**Cellular localization** Cytoplasm.


#### **Images**



Western blot detection of transfected COS-1 cells expressing STUB1/CHIP using ab2917.



Western blot analysis of STUB1/CHIP was performed by loading 20ug of total protein extracted from the left hemisphere (Left lane) or right hemisphere (right lane) of a normal C57BL/6 mouse brain per well on an SDS-PAGE gel. Proteins were transferred to a membrane, blocked with 5% non-fat dry milkand probed with a STUB1/CHIP polyclonal antibody (ab2915) at a dilution of 1:1000, followed by a HRP-conjugated goat anti-rabbit lgG secondary antibody. Detection was performed using a chemiluminescent substrate.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.com/abpromise">https://www.abcam.com/abpromise</a> or contact our technical team.

### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors