abcam

Product datasheet

Anti-Transketolase antibody ab233179

9 Images

Overview

Product name Anti-Transketolase antibody

Description Rabbit polyclonal to Transketolase

Host species Rabbit

Tested applications Suitable for: WB, IHC-P

Species reactivity Reacts with: Human

Predicted to work with: Mouse, Rat, Cow, Chimpanzee, Cynomolgus monkey, Orangutan

A

Immunogen Recombinant fragment (His-T7-tag) corresponding to Human Transketolase aa 1-300. Expressed

in E.coli. N-terminal tags. Database link: <u>P29401</u>

Run BLAST with
Run BLAST with

Positive control IHC-P: Human breast cancer, glioma, cerebrum, liver cancer, lung cancer, kidney and thyroid

tissues. WB: A549 cell lysate; Recombinant human Transketolase protein.

General notesThe Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

Storage buffer pH: 7.40

Preservative: 0.011% Proclin 300

Constituents: 55.77% Glycerol (glycerin, glycerine), 44.219% PBS

Purity Protein A purified

Purification notes ab233179 was purified by antigen-specific affinity chromatography followed by Protein A affinity

1

chromatography.

Clonality Polyclonal

Isotype IgG

Applications

The Abpromise guarantee Our Abpromise guarantee covers the use of ab233179 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
WB		Use a concentration of 0.2 - 2 µg/ml. Predicted molecular weight: 68 kDa.
IHC-P		Use a concentration of 5 - 20 µg/ml.

Target

Sequence similarities

Belongs to the transketolase family.

Images




Anti-Transketolase antibody (ab233179) at 1 μ g/ml + A549 (human lung carcinoma cell line) cell lysate

Secondary

HRP-Linked Goat Anti-Rabbit IgG Polyclonal at 0.2 µg/ml


Predicted band size: 68 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Transketolase antibody (ab233179)

Formalin-fixed, paraffin-embedded human thyroid tissue stained for Transketolase using ab233179 at 10 $\mu g/ml$ in immunohistochemical analysis.


Secondary used: HRP-Linked Goat Anti-Rabbit lgG Polyclonal at 2 µg/ml. DAB staining.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Transketolase antibody (ab233179)

Formalin-fixed, paraffin-embedded human breast cancer tissue stained for Transketolase using ab233179 at 10 μ g/ml in immunohistochemical analysis.


Secondary used: HRP-Linked Goat Anti-Rabbit lgG Polyclonal at 2 μ g/ml. DAB staining.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Transketolase antibody (ab233179)

Formalin-fixed, paraffin-embedded human glioma tissue stained for Transketolase using ab233179 at 10 $\mu g/ml$ in immunohistochemical analysis.


Secondary used: HRP-Linked Goat Anti-Rabbit lgG Polyclonal at 2 μ g/ml. DAB staining.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Transketolase antibody (ab233179)

Formalin-fixed, paraffin-embedded human cerebrum tissue stained for Transketolase using ab233179 at 10 μ g/ml in immunohistochemical analysis.


Secondary used: HRP-Linked Goat Anti-Rabbit lgG Polyclonal at 2 μ g/ml. DAB staining.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Transketolase antibody (ab233179)

Formalin-fixed, paraffin-embedded human liver cancer tissue stained for Transketolase using ab233179 at 10 $\mu g/ml$ in immunohistochemical analysis.


Secondary used: HRP-Linked Goat Anti-Rabbit $\lg G$ Polyclonal at 2 $\mu g/ml$. DAB staining.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Transketolase antibody (ab233179)

Formalin-fixed, paraffin-embedded human lung cancer tissue stained for Transketolase using ab233179 at 10 μ g/ml in immunohistochemical analysis.


Secondary used: HRP-Linked Goat Anti-Rabbit lgG Polyclonal at 2 μ g/ml. DAB staining.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Transketolase antibody (ab233179)

Formalin-fixed, paraffin-embedded human kidney tissue stained for Transketolase using ab233179 at 10 μ g/ml in immunohistochemical analysis.

Secondary used: HRP-Linked Goat Anti-Rabbit lgG Polyclonal at 2 µg/ml. DAB staining.



Anti-Transketolase antibody (ab233179) at 2 μ g/ml + Recombinant human Transketolase protein

Predicted band size: 68 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

• Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors