abcam

Product datasheet

Anti-TTI2 antibody - C-terminal ab176698

2 Images

Overview

Product name Anti-TTI2 antibody - C-terminal

Description Rabbit polyclonal to TTI2 - C-terminal

Host species Rabbit

Tested applications

Suitable for: WB, IP

Species reactivity

Reacts with: Human

Immunogen Synthetic peptide within Human TTI2 aa 458-508 (C terminal). The exact sequence is proprietary.

(NP_079391.2).

Sequence:

ATDCLILLDRCSQGRVKGLLAKIPQSCEDRKVVNYIRKVQQ

VSEGAPYN

Database link: **Q6NXR4**

Run BLAST with
Run BLAST with

Positive control HeLa, 293T and Jurkat whole cell lysates.

General notes

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

Storage buffer pH: 6.8

Preservative: 0.09% Sodium azide

Constituents: 0.1% BSA, 99% Tris buffered saline

Purity Immunogen affinity purified

1

Clonality

Polyclonal

Isotype

lgG

Applications

The Abpromise guarantee


Our **Abpromise guarantee** covers the use of ab176698 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/2000 - 1/10000. Predicted molecular weight: 57 kDa.
IP		Use at 2-10 μg/mg of lysate.

Target

Images



Western blot - Anti-TTI2 antibody - C-terminal (ab176698)

All lanes: Anti-TTl2 antibody - C-terminal (ab176698) at 0.04

µg/ml

Lane 1 : HeLa whole cell lysate at 50 μg

Lane 2 : HeLa whole cell lysate at 15 μg


Lane 3: 293T whole cell lysate at 50 µg

Lane 4: Jurkat whole cell lysate at 50 µg

Developed using the ECL technique.

Predicted band size: 57 kDa

Exposure time: 3 minutes



Immunoprecipitation. ab176698 at 1 μ g/ml staining TTl2 in HeLa whole cell lysate immunoprecipitated using ab176698 at 6 μ g/mg lysate (1 mg/IP; 20% of IP loaded/lane). Lane 2: control lgG.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

(ab176698)

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors