

Product datasheet

Anti-VPS4B/MIG1 antibody [EPR9225] ab137027

Recombinant RabMAb

[2 References](#) [3 Images](#)

Overview

| | |
|----------------------------|---|
| Product name | Anti-VPS4B/MIG1 antibody [EPR9225] |
| Description | Rabbit monoclonal [EPR9225] to VPS4B/MIG1 |
| Host species | Rabbit |
| Tested applications | Suitable for: Flow Cyt (Intra), WB Unsuitable for: ICC/IF, IHC-P or IP |
| Species reactivity | Reacts with: Human |
| Immunogen | Synthetic peptide within Human VPS4B/MIG1 aa 350-450. The exact sequence is proprietary. |
| Positive control | HeLa, 293T, Jurkat, and HepG2 whole cell lysate (ab7900); 293T cells |
| General notes | <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p> |

Properties

| | |
|-----------------------------|--|
| Form | Liquid |
| Storage instructions | Shipped at 4°C. Store at -20°C. |
| Storage buffer | pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 9% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA, 50% Tissue culture supernatant |
| Purity | Tissue culture supernatant |
| Clonality | Monoclonal |

Clone number EPR9225
Isotype IgG

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab137027 in the following tested applications.
The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes |
|------------------|-----------|---|
| Flow Cyt (Intra) | | 1/10 - 1/100. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody. |
| WB | | 1/1000 - 1/10000. Predicted molecular weight: 49 kDa. |

Application notes Is unsuitable for ICC/IF, IHC-P or IP.

Target

Function Involved in late steps of the endosomal multivesicular bodies (MVB) pathway. Recognizes membrane-associated ESCRT-III assemblies and catalyzes their disassembly, possibly in combination with membrane fission. Redistributes the ESCRT-III components to the cytoplasm for further rounds of MVB sorting. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. In conjunction with the ESCRT machinery also appears to function in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis and enveloped virus budding (HIV-1 and other lentiviruses).

Tissue specificity Ubiquitously expressed.


Sequence similarities Belongs to the AAA ATPase family.
Contains 1 MIT domain.

Domain The MIT domain serves as an adapter for ESCRT-III proteins. It forms an asymmetric three-helix bundle that binds amphipathic MIM (MIT interacting motif) helices along the groove between MIT helices 2 and 3 present in a subset of ESCRT-III proteins thus establishing the canonical MIM-MIT interaction. In an extended conformation along the groove between helices 1 and 3, also binds to a type-2 MIT interacting motif (MIM2).

Post-translational modifications Phosphorylated upon DNA damage, probably by ATM or ATR.

Cellular localization Prevacuolar compartment membrane. Late endosome membrane. Membrane-associated in the prevacuolar endosomal compartment. Localized in HIV-1 particles purified from acutely infected cells.

Images



Western blot - Anti-VPS4B/MIG1 antibody [EPR9225] (ab137027)

All lanes : Anti-VPS4B/MIG1 antibody [EPR9225] (ab137027) at 1/1000 dilution

Lane 1 : HeLa cell lysate

Lane 2 : 293T cell lysate

Lane 3 : Jurkat cell lysate


Lane 4 : HepG2 cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 49 kDa



Flow Cytometry (Intracellular) - Anti-VPS4B/MIG1 antibody [EPR9225] (ab137027)

Intracellular flow cytometric analysis of permeabilized 293T cells labelling VPS4B/MIG1 with ab137027 at 1/10 dilution (red). Rabbit IgG negative control is green.

Why choose a recombinant antibody?

| | |
|---|---|
|  Research with confidence Consistent and reproducible results |  Long-term and scalable supply Recombinant technology |
|  Success from the first experiment Confirmed specificity |  Ethical standards compliant Animal-free production |

Anti-VPS4B/MIG1 antibody [EPR9225] (ab137027)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors