

Product datasheet

Anti-WIF1 antibody [MM0602-6D27] α b89935

★★★★★ [1 Abreviews](#) [1 References](#)

Overview

| | |
|----------------------------|---|
| Product name | Anti-WIF1 antibody [MM0602-6D27] |
| Description | Mouse monoclonal [MM0602-6D27] to WIF1 |
| Host species | Mouse |
| Tested applications | Suitable for: IHC-P, WB |
| Species reactivity | Reacts with: Human |
| Immunogen | Recombinant full length protein Human WIF1 |
| General notes | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> |

Properties

| | |
|-----------------------------|--|
| Form | Liquid |
| Storage instructions | Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles. |
| Storage buffer | Constituent: PBS |
| Purity | Protein G purified |
| Purification notes | The IgG fraction of culture supernatant was purified by Protein G affinity chromatography and filtered through a 0.2 μ m filter. |
| Clonality | Monoclonal |
| Clone number | MM0602-6D27 |
| Isotype | IgG1 |

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab89935 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes |
|-------------|-----------|--|
| IHC-P | ★★★★★ (1) | Use at an assay dependent concentration. |
| WB | | Use at an assay dependent concentration. |

Target

| | |
|------------------------------|--|
| Function | Binds to WNT proteins and inhibits their activities. May be involved in mesoderm segmentation. |
| Sequence similarities | Contains 5 EGF-like domains. Contains 1 WIF domain. |
| Cellular localization | Secreted. |

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors