

Product datasheet

Anti-Wilms Tumor Protein antibody [WT1/857] ab220212

★ ★ ★ ★ ★ [1 Abreviews](#) [2 References](#) [4 Images](#)

Overview

| | |
|----------------------------|---|
| Product name | Anti-Wilms Tumor Protein antibody [WT1/857] |
| Description | Mouse monoclonal [WT1/857] to Wilms Tumor Protein |
| Host species | Mouse |
| Tested applications | Suitable for: IHC-P |
| Species reactivity | Reacts with: Rat, Human |
| Immunogen | Recombinant fragment corresponding to Human Wilms Tumor Protein. Database link: P19544 |
| Positive control | IHC-P: Rat Testis, Human Fetal Kidney, Testis, and Mesothelioma. |
| General notes | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> |

Properties

| | |
|-----------------------------|---|
| Form | Liquid |
| Storage instructions | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle. |
| Storage buffer | pH: 7.2 Preservative: 0.05% Sodium azide Constituents: 0.05% BSA, PBS |
| Purity | Protein G purified |
| Clonality | Monoclonal |
| Clone number | WT1/857 |
| Isotype | IgG1 |
| Light chain type | kappa |

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab220212 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes |
|-------------|-----------|--|
| IHC-P | | Use a concentration of 0.5 - 1 µg/ml. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. |

Target

Function

Transcription factor that plays an important role in cellular development and cell survival. Regulates the expression of numerous target genes, including EPO. Plays an essential role for development of the urogenital system. Recognizes and binds to the DNA sequence 5'-CGCCCCCGC-3'. It has a tumor suppressor as well as an oncogenic role in tumor formation. Function may be isoform-specific: isoforms lacking the KTS motif may act as transcription factors. Isoforms containing the KTS motif may bind mRNA and play a role in mRNA metabolism or splicing. Isoform 1 has lower affinity for DNA, and can bind RNA.

Tissue specificity

Expressed in the kidney and a subset of hematopoietic cells.

Involvement in disease

Defects in WT1 are the cause of Frasier syndrome (FS) [MIM:136680]. FS is characterized by a slowly progressing nephropathy leading to renal failure in adolescence or early adulthood, male pseudohermaphroditism, and no Wilms tumor. As for histological findings of the kidneys, focal glomerular sclerosis is often observed. There is phenotypic overlap with Denys-Drash syndrome. Inheritance is autosomal dominant.

Defects in WT1 are the cause of Wilms tumor 1 (WT1) [MIM:194070]. WT is an embryonal malignancy of the kidney that affects approximately 1 in 10'000 infants and young children. It occurs both in sporadic and hereditary forms.

Defects in WT1 are the cause of Denys-Drash syndrome (DDS) [MIM:194080]. DDS is a typical nephropathy characterized by diffuse mesangial sclerosis, genital abnormalities, and/or Wilms tumor. There is phenotypic overlap with WAGR syndrome and Frasier syndrome. Inheritance is autosomal dominant, but most cases are sporadic.

Defects in WT1 are the cause of nephrotic syndrome type 4 (NPHS4) [MIM:256370]. A renal disease characterized clinically by proteinuria, hypoalbuminemia, hyperlipidemia and edema. Kidney biopsies show non-specific histologic changes such as focal segmental glomerulosclerosis and diffuse mesangial proliferation. Some affected individuals have an inherited steroid-resistant form and progress to end-stage renal failure. Most patients with NPHS4 show diffuse mesangial sclerosis on renal biopsy, which is a pathologic entity characterized by mesangial matrix expansion with no mesangial hypercellularity, hypertrophy of the podocytes, vacuolized podocytes, thickened basement membranes, and diminished patency of the capillary lumen.

Defects in WT1 are a cause of Meacham syndrome (MEACHS) [MIM:608978]. Meacham syndrome is a rare sporadically occurring multiple malformation syndrome characterized by male pseudohermaphroditism with abnormal internal female genitalia comprising a uterus and double or septate vagina, complex congenital heart defect and diaphragmatic abnormalities.


Note=A chromosomal aberration involving WT1 may be a cause of desmoplastic small round cell tumor (DSRCT). Translocation t(11;22)(p13;q12) with EWSR1.

Sequence similarities

Belongs to the EGR C2H2-type zinc-finger protein family.
Contains 4 C2H2-type zinc fingers.


Cellular localization

Nucleus. Cytoplasm. Shuttles between nucleus and cytoplasm; Nucleus > nucleoplasm and Nucleus speckle.

Images


Immunohistochemical analysis of formalin-fixed and paraffin-embedded rat testis tissue labeling Wilms Tumor protein with ab220212 at 1 µg/mL.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Wilms Tumor Protein antibody [WT1/857] (ab220212)




Immunohistochemical analysis of formalin-fixed and paraffin-embedded human Mesothelioma tissue labeling Wilms Tumor protein with ab220212 at 1 µg/mL.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Wilms Tumor Protein antibody [WT1/857] (ab220212)



Immunohistochemical analysis of formalin-fixed and paraffin-embedded human testis tissue labeling Wilms Tumor protein with ab220212 at 1 µg/mL.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Wilms Tumor Protein antibody [WT1/857] (ab220212)



Immunohistochemical analysis of formalin-fixed and paraffin-embedded human fetal kidney tissue labeling Wilms Tumor protein with ab220212 at 1 µg/mL.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Wilms Tumor Protein antibody [WT1/857] (ab220212)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors