

## Product datasheet

# Human Hepsidin-25 peptide ab31875

★★★★☆ [1 Abreviews](#) [4 References](#) [1 Image](#)

### Description

---

<b>Product name</b>	Human Hepsidin-25 peptide
<b>Purity</b>	> 70 % HPLC. 70 - 90% by HPLC
<b>Accession</b>	<b><u>P81172</u></b>
<b>Animal free</b>	No
<b>Nature</b>	Synthetic
<b>Species</b>	Human

### Specifications

---

Our **Abpromise guarantee** covers the use of **ab31875** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>Applications</b>	Blocking - Blocking peptide for Anti-Hepsidin-25 antibody ( <b><u>ab30760</u></b> ) ELISA
<b>Form</b>	Lyophilized
<b>Additional notes</b>	Storage buffer: 10% 0.5M acetic acid, 90% TK buffer. pH 3.6 Solutions of ab31875, diluted 10 times, in TBST (80%) and PBS (10%) have a pH of 5.1

### Preparation and Storage

---

<b>Stability and Storage</b>	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles. Information available upon request.
<b>Reconstitution</b>	Reconstitute in 10% 0.5M acetic acid, 90% TK buffer. If the peptide doesn't dissolve try an organic solvent like DMSO, then dilute using water or buffer. Gentle warming and sonication can effectively aid peptide solubilisation

### General Info

---

<b>Function</b>	Liver-produced hormone that constitutes the main circulating regulator of iron absorption and
-----------------	---

distribution across tissues. Acts by promoting endocytosis and degradation of ferroportin, leading to the retention of iron in iron-exporting cells and decreased flow of iron into plasma. Controls the major flows of iron into plasma: absorption of dietary iron in the intestine, recycling of iron by macrophages, which phagocytose old erythrocytes and other cells, and mobilization of stored iron from hepatocytes (PubMed:22306005).

Has strong antimicrobial activity against E.coli ML35P N.cinerea and weaker against S.epidermidis, S.aureus and group b streptococcus bacteria. Active against the fungus C.albicans. No activity against P.aeruginosa (PubMed:11113131, PubMed:11034317).

**Tissue specificity**

Highest expression in liver and to a lesser extent in heart and brain. Low levels in lung, tonsils, salivary gland, trachea, prostate gland, adrenal gland and thyroid gland. Secreted into the urine.

**Involvement in disease**

Hemochromatosis 2B

**Sequence similarities**

Belongs to the hepcidin family.

**Cellular localization**

Secreted.

---

**Images**

**Choose the best for your experiment from thousands of proteins and peptides**

- Reinforce the validity of your results with **protein controls**
- Validate specific, reliable reagents with **blocking peptides**
- Keep working on targets without specific antibodies with **fusion-tagged proteins**
- Get consistent, reproducible results with premium **bioactive proteins**

Human Hepcidin-25 peptide (ab31875)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

---

**Our Abpromise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors