# abcam

### Product datasheet

## Native Human Serum Albumin protein (HRP) ab8032

**Description** 

Product name Native Human Serum Albumin protein (HRP)

Expression system Native

Protein length Full length protein

Animal free No

**Nature** Native

Species Human
Predicted molecular weight 69 kDa

**Conjugation** HRP

**Description** Native Human Human Serum Albumin protein (HRP)

#### **Specifications**

Our Abpromise guarantee covers the use of ab8032 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

**Applications** Western blot

Dot blot ELISA

Immunomicroscopy

Form Lyophilized

#### **Preparation and Storage**

**Stability and Storage** Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Preservative: 0.01% Gentamicin sulphate

 $Constituents: 1\% \ Polyethylene \ glycol, \ 0.42\% \ Potassium \ phosphate, \ 0.87\% \ Sodium \ chloride$ 

**Reconstitution** Restore with 1.0 mL of deionized water (or equivalent).

#### **General Info**

Function Serum albumin, the main protein of plasma, has a good binding capacity for water, Ca(2+), Na(+),

1

K(+), fatty acids, hormones, bilirubin and drugs. Its main function is the regulation of the colloidal osmotic pressure of blood. Major zinc transporter in plasma, typically binds about 80% of all

plasma zinc.

**Tissue specificity** Plasma.

Involvement in disease Defects in ALB are a cause of familial dysalbuminemic hyperthyroxinemia (FDH) [MIM:103600].

 ${\sf FDH}\ is\ a\ form\ of\ euthyroid\ hyperthyroxinemia\ that\ is\ due\ to\ increased\ affinity\ of\ ALB\ for\ T(4).\ It\ is$ 

the most common cause of inherited euthyroid hyperthyroxinemia in Caucasian population.

**Sequence similarities**Belongs to the ALB/AFP/VDB family.

Contains 3 albumin domains.

Post-translational

modifications

Kenitra variant is partially O-glycosylated at Thr-620. It has two new disulfide bonds Cys-600 to

Cys-602 and Cys-601 to Cys-606.

Glycated in diabetic patients.

Phosphorylation sites are present in the extracelllular medium.

Acetylated on Lys-223 by acetylsalicylic acid.

Cellular localization Secreted.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.com/abpromise">https://www.abcam.com/abpromise</a> or contact our technical team.

#### Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors