

# Native Mouse Vitronectin/S-Protein ab92727

★★★★★ [1 Abreviews](#) [4 References](#) [1 Image](#)

### Description

<b>Product name</b>	Native Mouse Vitronectin/S-Protein
<b>Purity</b>	> 95 % SDS-PAGE. ab92727 was prepared from fresh Mouse plasma using urea as a denaturant.
<b>Expression system</b>	Native
<b>Protein length</b>	Full length protein
<b>Animal free</b>	No
<b>Nature</b>	Native
<b>Species</b>	Mouse

### Specifications

Our **Abpromise guarantee** covers the use of **ab92727** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>Applications</b>	Functional Studies SDS-PAGE
<b>Form</b>	Liquid

### Preparation and Storage

<b>Stability and Storage</b>	Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles. pH: 7.40 Constituents: 0.82% Sodium phosphate, 0.58% Sodium chloride
------------------------------	--

### General Info

<b>Function</b>	Vitronectin is a cell adhesion and spreading factor found in serum and tissues. Vitronectin interact with glycosaminoglycans and proteoglycans. Is recognized by certain members of the integrin family and serves as a cell-to-substrate adhesion molecule. Inhibitor of the membrane-damaging effect of the terminal cytolytic complement pathway. Somatomedin-B is a growth hormone-dependent serum factor with protease-inhibiting activity.
-----------------	---

<b>Tissue specificity</b>	Plasma.
<b>Sequence similarities</b>	Contains 4 hemopexin repeats. Contains 1 SMB (somatomedin-B) domain.
<b>Domain</b>	The SMB domain mediates interaction with SERPINE1/PAI1. The heparin-binding domain mediates interaction with insulin.
<b>Post-translational modifications</b>	Sulfated on 2 tyrosine residues. N- and O-glycosylated. Phosphorylation on Thr-69 and Thr-76 favors cell adhesion and spreading. It has been suggested that the active SMB domain may be permitted considerable disulfide bond heterogeneity or variability, thus two alternate disulfide patterns based on 3D structures are described with 1 disulfide bond conserved in both. Phosphorylation sites are present in the extracellular medium.
<b>Cellular localization</b>	Secreted, extracellular space.

## Images



10% SDS-PAGE showing ab92727  
Lane 1 ab92727 Non-reduced (3µg)  
Lane 2 ab92727 Reduced (3µg)  
Lane 3 Molecular weight markers

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors