

Product datasheet

Recombinant Human ADAM10 protein ab82822

1 Image

Description

Product name	Recombinant Human ADAM10 protein
Purity	> 95 % SDS-PAGE.
Expression system	Escherichia coli
Accession	<u>O14672</u>
Protein length	Protein fragment
Animal free	No
Nature	Recombinant
Species	Human
Amino acids	214 to 672
Tags	His tag N-Terminus
Additional sequence information	The His tag is a 6-His tag.

Specifications

Our **Abpromise guarantee** covers the use of **ab82822** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	SDS-PAGE
Form	Liquid

Preparation and Storage

Stability and Storage	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. pH: 7.20 Constituents: PBS, 50% Glycerol
------------------------------	--

General Info

Function	Cleaves the membrane-bound precursor of TNF-alpha at '76-Ala-Val-77' to its mature soluble form. Responsible for the proteolytical release of soluble JAM3 from
-----------------	---

endothelial cells surface. Responsible for the proteolytic release of several other cell-surface proteins, including heparin-binding epidermal growth-like factor, ephrin-A2 and for constitutive and regulated alpha-secretase cleavage of amyloid precursor protein (APP). Contributes to the normal cleavage of the cellular prion protein. Involved in the cleavage of the adhesion molecule L1 at the cell surface and in released membrane vesicles, suggesting a vesicle-based protease activity. Controls also the proteolytic processing of Notch and mediates lateral inhibition during neurogenesis.

Tissue specificity

Expressed in spleen, lymph node, thymus, peripheral blood leukocyte, bone marrow, cartilage, chondrocytes and fetal liver.

Sequence similarities

Contains 1 disintegrin domain.
Contains 1 peptidase M12B domain.

Domain

The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.


Post-translational modifications

The precursor is cleaved by a furin endopeptidase.

Cellular localization

Cell membrane. Endomembrane system. Is localized in the plasma membrane but is predominantly expressed in the Golgi apparatus and in released membrane vesicles derived likely from the Golgi.

Images



ab82822 on SDS-PAGE.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors