

Recombinant Human C1D protein ab156979

1 Image

Description

Product name	Recombinant Human C1D protein
Purity	> 85 % SDS-PAGE. ab156979 is purified using conventional chromatography techniques.
Expression system	Escherichia coli
Accession	<u>Q13901</u>
Protein length	Full length protein
Animal free	No
Nature	Recombinant
Species	Human
Sequence	MGSSHHHHHH SSGLVPRGSH MGSMAGEEIN EDYPVEIHEYLSAFENSIGA VDEMLKTMMMS VSRNELLQKL DPLEQAKVDL VSAYTLNSMF WVYLATQGVN PKEHPVKQEL ERIRVYMNRV KEITDKKKAG KLDRGAASRF VKNALWEPKS KNASKVANKG KSKS
Predicted molecular weight	18 kDa including tags
Amino acids	1 to 141
Tags	His tag N-Terminus

Specifications

Our **Abpromise guarantee** covers the use of **ab156979** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	Mass Spectrometry SDS-PAGE
Mass spectrometry	MALDI-TOF
Form	Liquid

Preparation and Storage

Stability and Storage	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -
------------------------------	--

80°C. Avoid freeze / thaw cycle.


pH: 8.00

Constituents: 0.03% DTT, 0.32% Tris HCl, 50% Glycerol (glycerin, glycerine), 1.17% Sodium chloride

General Info

Function	Plays a role in the recruitment of the RNA exosome complex to pre-rRNA to mediate the 3'-5' end processing of the 5.8S rRNA; this function may include MPHOSPH6. Can activate PRKDC not only in the presence of linear DNA but also in the presence of supercoiled DNA. Can induce apoptosis in a p53/TP53 dependent manner. May regulate the TRAX/TSN complex formation. Potentiates transcriptional repression by NR1D1 and THRB.
Tissue specificity	Ubiquitous. Expressed at very high levels in the hippocampus, medulla oblongata, mammary gland, thyroid and salivary gland. Expressed at high levels in the fetal; lung, liver and kidney. Expressed at low levels in skeletal muscle, appendix, heart, lung and colon.
Sequence similarities	Belongs to the C1D family.
Post-translational modifications	Phosphorylated by PRKDC.
Cellular localization	Cytoplasm. Nucleus > nucleolus. EXOSC10 is required for nucleolar localization.

Images



15% SDS-PAGE analysis of ab156979 (3µg).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors