

Recombinant human CD116 protein (Fc Chimera)  
ab83993

2 Images

Description

Product name	Recombinant human CD116 protein (Fc Chimera)
Biological activity	ab83993 bound to protein A sepharose beads was able to pull down its ligand, GM-CSF.
Purity	> 95 % SDS-PAGE.
Expression system	HEK 293 cells
Accession	<b><u>P15509</u></b>
Protein length	Full length protein
Animal free	No
Nature	Recombinant
Species	Human

Sequence

Theoretical Sequence:  
EKSDLRTVAPASSLNVRFDSRTMNLSDCQENTTFSKCF  
LTDKKNRVV  
EPRLSNNECSCTFREICLHEGVTFEVHVNTSQRGFQKLL  
YPNSGREG  
TAAQNFSCFIYNADLMNCTWARGPTAPRDVQYFLYRNSK  
RRREIRCP  
YYIQDSGTHVGCHLDNLSGLTSRNYFLVNGTSREIGIQFFD  
SLLDTKK  
IERFNPPSNVTVRCNTTHCLVRWKQPRTYQKLSYLDFQYQ  
LDVHRKNT  
QPGTENLLINVSGDLENRYNFPSSSEPRAKHSVKIRAADVRI  
LNWSSWS  
EAIEFGSDDGGSSNTKVDDKVEPKSCDKTHTCPPCPAPE  
LLGGPSVFL  
FPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDG  
VEVHNAKTK  
PREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALP  
APIEKTISK  
AKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIA  
VEWESNGQ  
PENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSC

**Additional sequence information** Fusion of aa 1-320 of human GM-CSF Receptor alpha and aa 90-330 of Fc region of human IgG1 (P01857). The chimeric protein was expressed in modified human 293 cells.

## Specifications

---

Our **Abpromise guarantee** covers the use of **ab83993** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>Applications</b>	SDS-PAGE
<b>Form</b>	Lyophilized
<b>Additional notes</b>	ab83993 bound to protein A sepharose beads was able to pull down its ligand, GM-CSF. This product was previously labelled as GM-CSF Receptor alpha

## Preparation and Storage

---

<b>Stability and Storage</b>	Shipped at 4°C. Store at +4°C.  Constituents: 1% Human serum albumin, 10% Trehalose  This product is an active protein and may elicit a biological response in vivo, handle with caution.
<b>Reconstitution</b>	It is recommended that 0.5 ml of sterile phosphate-buffered saline be added to the vial. Following reconstitution short-term storage at 4°C is recommended, and longer-term storage of aliquots at -18 to -20°C. Repeated freeze thawing is not recommended.

## General Info


---

<b>Function</b>	Low affinity receptor for granulocyte-macrophage colony-stimulating factor. Transduces a signal that results in the proliferation, differentiation, and functional activation of hematopoietic cells.
<b>Involvement in disease</b>	Defects in CSF2RA are the cause of pulmonary surfactant metabolism dysfunction type 4 (SMDP4) [MIM:300770]. A rare lung disorder due to impaired surfactant homeostasis. It is characterized by alveolar filling with floccular material that stains positive using the periodic acid-Schiff method and is derived from surfactant phospholipids and protein components. Excessive lipoproteins accumulation in the alveoli results in severe respiratory distress.
<b>Sequence similarities</b>	Belongs to the type I cytokine receptor family. Type 5 subfamily.
<b>Domain</b>	The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding. The box 1 motif is required for JAK interaction and/or activation.
<b>Cellular localization</b>	Secreted and Cell membrane.

---

## Images

---




Functional Studies - Recombinant human CD116 protein (Fc Chimera) (ab83993)

Densitometry of protein isoforms visualised by 2-DE.

The densitometry scan demonstrates the purified human cell expressed protein exists in multiple isoforms, which differ according to their level of post-translational modification.

The triangle indicates the theoretical MW and pI of the protein.



SDS-PAGE - Recombinant human CD116 protein (Fc Chimera) (ab83993)

1D SDS-PAGE of ab83993 before and after treatment with glycosidases to remove oligosaccharides.

Lane 1: ab83993

Lane 2: ab83993 treated with PNGase F to remove potential N-linked glycans

Lane 3: ab83993 treated with a glycosidase cocktail to remove potential N- and O-linked glycans.

Approximately 5 µg of protein was loaded per lane; Gel was stained using Deep Purple™.

Drop in MW after treatment with PNGase F indicates presence of N-linked glycans. A tightening of the band after treatment with the glycosidase cocktail indicates O-linked glycans may be present. Additional bands in lane 2 and lane 3 are glycosidase enzymes.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

- 
- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors