abcam

Product datasheet

Recombinant Human EBP1 protein ab95350

2 Images

Description

Product name Recombinant Human EBP1 protein

Purity > 95 % SDS-PAGE.

ab95350 is purified using conventional chromatography techniques.

Expression system Escherichia coli

Protein length Full length protein

Animal free No

Nature Recombinant

Species Human

Sequence MSGEDEQQEQ TIAEDLVVTK YKMGGDIANR

VLRSLVEASS SGVSVLSLCE KGDAMIMEET

GKIFKKEKEM KKGIAFPTSI SVNNCVCHFS PLKSDQDYIL

KEGDLVKIDL GVHVDGFIAN VAHTFVVDVA
QGTQVTGRKA DVIKAAHLCA EAALRLVKPG
NQNTQVTEAW NKVAHSFNCT PIEGMLSHQL
KQHVIDGEKT IIQNPTDQQK KDHEKAEFEV
HEVYAVDVLV SSGEGKAKDA GQRTTIYKRD
PSKQYGLKMK TSRAFFSEVE RRFDAMPFTL
RAFEDEKKAR MGVVECAKHE LLQPFNVLYE
KEGEFVAQFK FTVLLMPNGP MRITSGPFEP
DLYKSEMEVQ DAELKALLQS SASRKTQKKK
KKKASKTAEN ATSGETLEEN EAGDLEHHHH HH

Specifications

Our Abpromise guarantee covers the use of ab95350 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications Western blot

SDS-PAGE

Form Liquid

Preparation and Storage

1

Stability and Storage

Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

pH: 8.00

Constituents: 0.316% Tris HCI, 10% Glycerol (glycerin, glycerine)

General Info

Function

May play a role in a ERBB3-regulated signal transduction pathway. Seems be involved in growth regulation. Acts a corepressor of the androgen receptor (AR) and is regulated by the ERBB3 ligand neuregulin-1/heregulin (HRG). Inhibits transcription of some E2F1-regulated promoters, probably by recruiting histone acetylase (HAT) activity. Binds RNA. Associates with 28S, 18S and 5.8S mature rRNAs, several rRNA precursors and probably U3 small nucleolar RNA. May be involved in regulation of intermediate and late steps of rRNA processing. May be involved in ribosome assembly. Mediates cap-independent translation of specific viral IRESs (internal ribosomal entry site).

Tissue specificity

Expressed in several cell lines tested, including primary and transformed cell lines.

Sequence similarities

Belongs to the peptidase M24 family.

Post-translational modifications


Phosphorylated on serine and threonine residues. Phosphorylation is enhanced by HRG treatment. Basal phosphorylation is PKC-dependent and HRG-induced phosphorylation is

predominantly PKC-independent.

Cellular localization


Cytoplasm. Nucleus > nucleolus. Tranlocates to the nucleus upon treatment with HRG.

Images



SDS-PAGE - Recombinant Human EBP1 protein (ab95350)

15% SDS-PAGE showing ab95350 at approximately 45 kDa (3 μ g).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors