# abcam

## Product datasheet

## Recombinant human IGF1 protein (Active) ab270062

\*\*\* \* \* 1 Abreviews 5 References 4 Images

**Description** 

Product name Recombinant human IGF1 protein (Active)

**Biological activity** Fully biologically active as determined by dose-depended proliferation of MCF-7 cells.

ED<sub>50</sub> is  $\leq$  0.9 ng/mL, corresponding to a specific activity of 1.11 x 10<sup>6</sup> units/mg.

Purity >= 95 % SDS-PAGE.

Endotoxin level< 0.005 Eu/μg</th>Expression systemHEK 293 cells

Accession P05019

Protein length Full length protein

Animal free Yes
Carrier free Yes

Nature Recombinant

**Species** Human

**Sequence** GP ETLCGAELVD ALQFVCGDRG FYFNKPTGYG

SSSRRAPQTG IVDECCFRSC DLRRLEMYCA PLKPAKSA

Predicted molecular weight 8 kDa

Molecular weight information M - 2.84 Da (Calculated mass 7711.84 Da) GP ETLCGAELVD ALQFVCGDRG FYFNKPTGYG

SSSRRAPQTG IVDECCFRSC DLRRLEMYCA PLKPAKSA

Amino acids 49 to 118

Additional sequence information N-terminal glycine Full-length mature chain lacking the signal peptide and both pro-peptides.

**Specifications** 

Our Abpromise guarantee covers the use of ab270062 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

**Applications** Cell Culture

SDS-PAGE

**HPLC** 

Mass Spectrometry

**Functional Studies** 

Form Lyophilized

Additional notes This protein is filter sterilized prior to aliquoting and lyophilization. All aliquoting and lyophilization

steps are performed in a sterile environment

#### **Preparation and Storage**

**Stability and Storage** Shipped at Room Temperature. Store at Room Temperature.

pH: 6.00

Constituents: 0.727% Dibasic monohydrogen potassium phosphate, 0.248% Monobasic

dihydrogen potassium phosphate, 10.26% Trehalose

Buffer lyophilized from.

This product is an active protein and may elicit a biological response in vivo, handle with caution.

**Reconstitution** Reconstitute with phosphate buffered saline. Store lyophilized form at room temperature.

Reconstitute, aliquot and store at -80°C for 12 months or +4°C for 1 week. Avoid repeated freeze-

thaw. Lyophilized contents may appear as either a translucent film or a white powder. This

variance does not affect the quality of the product.

#### **General Info**

**Function** The insulin-like growth factors, isolated from plasma, are structurally and functionally related to

insulin but have a much higher growth-promoting activity. May be a physiological regulator of [1-14C]-2-deoxy-D-glucose (2DG) transport and glycogen synthesis in osteoblasts. Stimulates glucose transport in rat bone-derived osteoblastic (PyMS) cells and is effective at much lower concentrations than insulin, not only regarding glycogen and DNA synthesis but also with regard to

enhancing glucose uptake.

**Involvement in disease**Defects in IGF1 are the cause of insulin-like growth factor I deficiency (IGF1 deficiency)


[MIM:608747]. IGF1 deficiency is an autosomal recessive disorder characterized by growth

retardation, sensorineural deafness and mental retardation.

Sequence similarities Belongs to the insulin family.


Cellular localization Secreted.

## **Images**




Fully biologicaly active as determined by dose-depended proliferation of MCF-7 cells. The ED $_{50}$  is  $\leq 0.9$  ng/mL, corresponding to a specific activity of 1.11 x  $10^6$  units/mg.

Functional Studies - Recombinant human IGF1 protein (Active) (ab270062)




SDS-PAGE analysis of ab270062.

SDS-PAGE - Recombinant human IGF1 protein (Active) (ab270062)



### Purity ≥95%

The spectrum was recorded using a 1260 Infinity II HPLC system with DAD and a MabPac RP column (3.0x100 mm, 4  $\mu m$ ). 5  $\mu L$  of purified protein was injected and the gradient run from 80 % water:TFA (99.9:0.1 v/v) and 20 % acetonitrile:water:TFA (90:9.9:0.1 v/v/v) to 20 % water:TFA (99.9:0.1 v/v) and 80 % acetonitrile:water:TFA (90:9.9:0.1 v/v/v) within 3 minutes followed by an isocratic step for another 3 min. Flow rate was 0.5 mL/min and the column compartment temperature was 50 °C.



## M - 2.84 Da (Calculated mass 7711.84 Da)

The spectrum was recorded with a 6545XT AdvanceBio LC/Q-TOF (Agilent Technologies) and a MabPac RP column (42.1x50 mm, 4  $\mu$ m, Thermo Scientific). 5  $\mu$ L of purified protein was injected and the gradient run from 85 % water:FA (99.9:0.1 v/v) and 15 % acetonitrile:FA (90:9.9:0.1 v/v/v) to 55 % water:FA (99.9:0.1 v/v) and 45 % acetonitrile:FA (90:9.9:0.1 v/v/v) within 3 minutes followed by an isocratic step for another 2.5 min. Flow rate was 0.4 mL/min and the column compartment temperature was 60 °C. Data was analysed and deconvoluted using the Bioconfirm software (Agilent Technologies).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.com/abpromise">https://www.abcam.com/abpromise</a> or contact our technical team.

## Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors