

Recombinant human Neuropilin 1 protein (Fc Chimera Active) ab221023

3 Images

Description

Product name	Recombinant human Neuropilin 1 protein (Fc Chimera Active)		
Biological activity	Measured by its binding ability in a functional ELISA. Immobilized ab221023 at 2 µg/mL (100 µL/well) can bind Biotinylated Human VEGF165 with a linear range of 5-40 ng/mL.		
Purity	> 95 % SDS-PAGE.		
Endotoxin level	< 1.000 Eu/µg		
Expression system	HEK 293 cells		
Accession	<u>O14786-2</u>		
Protein length	Full length protein		
Animal free	No		
Nature	Recombinant		
Species	Human		
Sequence	FRNDKCGDT IKIESPGYLT SPGYPHSYHP SEKCEWLIQA PDPYQRIMIN FNPFDLEDR DCKYDYVEVF DGENENGHFR GKFCGKIAPP PVVSSGPFLF IKFVSDYETH GAGFSIRYEI FKRGPESQSN YTPSGVIKS PGFPEKYPNS LECTYMFVP KMSEIILEFE SFDLEPDSNP PGGMFCRYDR LEWDGFDPDV GPHIGRYCGQ KTPGRIRSSS GILSMVFYTD SAIKEGFSA NYSVLQSSVS EDFKCMEALG MESGEIHSDQ ITASSQYSTN WSAERSRLNY PENGWTPGED SYREWIQVDL GLLRFVTAVG TQGAISKETK KKYVKTYKI DVSSNGEDWI TIKEGNKPVL FQGNTNPTDV VVAVFPKPLI TRFVRIKPAT WETGISMRFE VYGCKITDYP CSGMLGMVSG LISDSQITSS NQGDRNWMPE NIRLVTSRSG WALPPAPHSY INEWLQIDLG EEKIVRGIII QGGKHRENVK FMRKFKIGYS NNGSDWKMM DDSKRKAKSF EGNNNYDTPE LRTFPALSTR FIRIYPERAT HGGLGLRMEL LGCEVEAPTA GPTTPNGNLV DECDDDQANC HSGTGDDDFQL TGGTTVLATE KPTVIDSTIQ SGIK		
Predicted molecular weight	97 kDa		

Amino acids	22 to 644
Additional sequence information	This product is for the mature full length protein. The signal peptide is not included. Fused with a human IgG1 Fc tag (Pro 100–Lys 330; P01857) at the C-terminus (AAH07533).

Specifications

Our **Abpromise guarantee** covers the use of **ab221023** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	SDS-PAGE Functional Studies
Form	Lyophilized


Preparation and Storage

Stability and Storage	Shipped at 4°C. Store at -20°C or -80°C. Avoid freeze / thaw cycle. pH: 7.4 Constituents: 0.61% Tris, 0.75% Glycine, 5% Trehalose, Sodium chloride, L-Arginine Lyophilized from 0.22 µm filtered solution. This product is an active protein and may elicit a biological response in vivo, handle with caution.
Reconstitution	Reconstitute with sterile deionized water to a concentration of 200 µg/ml.

General Info


Function	The membrane-bound isoform 1 is a receptor involved in the development of the cardiovascular system, in angiogenesis, in the formation of certain neuronal circuits and in organogenesis outside the nervous system. It mediates the chemorepulsant activity of semaphorins. It binds to semaphorin 3A, The PLGF-2 isoform of PGF, The VEGF-165 isoform of VEGF and VEGF-B. Coexpression with KDR results in increased VEGF-165 binding to KDR as well as increased chemotaxis. It may regulate VEGF-induced angiogenesis. The soluble isoform 2 binds VEGF-165 and appears to inhibit its binding to cells. It may also induce apoptosis by sequestering VEGF-165. May bind as well various members of the semaphorin family. Its expression has an averse effect on blood vessel number and integrity.
Tissue specificity	The expression of isoforms 1 and 2 does not seem to overlap. Isoform 1 is expressed by the blood vessels of different tissues. In the developing embryo it is found predominantly in the nervous system. In adult tissues, it is highly expressed in heart and placenta; moderately in lung, liver, skeletal muscle, kidney and pancreas; and low in adult brain. Isoform 2 is found in liver hepatocytes, kidney distal and proximal tubules.
Sequence similarities	Belongs to the neuropilin family. Contains 2 CUB domains. Contains 2 F5/8 type C domains. Contains 1 MAM domain.
Cellular localization	Secreted and Cell membrane.

Images




Functional Studies - Recombinant human Neuropilin 1 protein (Fc Chimera Active) (ab221023)

Immobilized Human VEGF165 (Tag Free) at 2 µg/mL (100 µL/well) can bind Human NRP1 (Fc Tag) with a linear range of 0.2-3 ng/mL (QC tested).



SDS-PAGE - Recombinant human Neuropilin 1 protein (Fc Chimera Active) (ab221023)

SDS-PAGE analysis of ab221023 under reducing conditions, stained overnight with Coomassie Blue. As a result of glycosylation, the reduced protein migrates as 120-125 kDa.



Functional Studies - Recombinant human Neuropilin 1 protein (Fc Chimera Active) (ab221023)

Immobilized ab221023 at 2 µg/mL (100 µL/well) can bind Biotinylated Human VEGF165 with a linear range of 5-40 ng/mL.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors