

## Product datasheet

# Recombinant Human NFkB p105 / p50 protein ab114185

[1 Image](#)

### Description

---

|                          |   |
|--------------------------|---|
| <b>Product name</b>      | Recombinant Human NFkB p105 / p50 protein |
| <b>Expression system</b> | Wheat germ                                |
| <b>Accession</b>         | <b><u>P19838</u></b>                      |
| <b>Protein length</b>    | Full length protein                       |
| <b>Animal free</b>       | No  |
| <b>Nature</b>            | Recombinant                               |
| <b>Species</b>           | Human                                     |

|                 |  |
|-----------------|--|
| <b>Sequence</b> | MAEDDPYLGRPEQMFHLDPSLTHITFNPEVFQPQMALPT<br>ADGPYLQILEQ<br>PKQRGFRFRYVCEGPSHGGLPGASSEKNKKSYPQVKICN<br>YVGPAAKVMVQL<br>VTNGKNIHLHAHSLVGKHCEDGICTVTAGPKDMVVGAFANL<br>GILHVTKKKV<br>FETLEARMTEACIRGYNPGLLVHPDLAYLQAEGGGDRQLG<br>DREKELIRQA<br>ALQQTKEMDLSVVRMLMFTAFLPDSTGSFTRRLEPVVSDAI<br>YDSKAPNASN<br>LKMVRMDRTAGCVTGGEEMLLCDKVQKDDIQIRFYEEEEEN<br>GGVWEGFGD<br>FSPTDVHRQFAIVFKTPKYKDINITKPASVFVQLRRKSDLET<br>SEPKPFLY<br>YPEIKDKKEEVQRKRQKLMFNFSDSFGGGSGAGAGGGGM<br>FGSGGGGGGTGS<br>TGPGYSFPHYGFPTYGGITFHPGTTKSNAGMKHGTMDTES<br>KKDPEGCDKS<br>DDKNTVNLFGKVIETTEQDQEPSEATVGNGEVTLTYATGT<br>KEESAGVQDN<br>LFLEKAMQLAKRHANALFDYAVTGDVKMLLAVQRHLTAV<br>QDENGDSVLHL<br>AIIHLHSQLVRLDLEVTSGSLISDDIINMRNDLYQTPLHLAVITK<br>QEDVVE<br>DLLRAGADLSLLDRLGNSVLHLAAKEGHDKVLSILLKHKK<br>AALLDHPNG |
|-----------------|--|

DGLNAIHLAMMSNSLPCLLLLVAAGADVNAQEKGSRGTA  
LHLAVEHDNIS  
LAGCLLLEGDAHVDSTTYDGTTPHIAAGRGSTRLAALLKA  
AGADPLVEN  
FEPLYDLDDSWENAGEDEGVVPGTTPLDMATSWQVFDIL  
NGKPYEPEFTS  
DDLLAQGDMKQLAEDVKLQLYKLEIPDPDKNWATLAQK  
LGLGILNNAFR  
LSPAPSKTLMNDNYEVSGGTVRELVEALRQMGYTEAIEVIQ  
AASSPVKTTS  
QAHSLPLSPASTRQQIDELRSDSVCDSGVETSFRKLSFT  
ESLTSGASLL TLNKMPHDYQGEGPLEGKI

**Predicted molecular weight** 133 kDa including tags  
**Amino acids** 1 to 968

## Specifications

---

Our **Abpromise guarantee** covers the use of **ab114185** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

**Applications** Western blot  
ELISA  
SDS-PAGE

**Form** Liquid

## Preparation and Storage

---

**Stability and Storage** Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles.  
pH: 8.00  
Constituents: 0.3% Glutathione, 0.79% Tris HCl

## General Info

---

**Function** NF-kappa-B is a pleiotropic transcription factor which is present in almost all cell types and is involved in many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52 and the heterodimeric p65-p50 complex appears to be most abundant one. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-kappa-B inhibitor (I-kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. NF-kappa-B heterodimeric p65-p50 and RelB-p50 complexes are transcriptional

activators. The NF-kappa-B p50-p50 homodimer is a transcriptional repressor, but can act as a transcriptional activator when associated with BCL3. NFKB1 appears to have dual functions such as cytoplasmic retention of attached NF-kappa-B proteins by p105 and generation of p50 by a cotranslational processing. The proteasome-mediated process ensures the production of both p50 and p105 and preserves their independent function, although processing of NFKB1/p105 also appears to occur post-translationally. p50 binds to the kappa-B consensus sequence 5'-GGRNNYCC-3', located in the enhancer region of genes involved in immune response and acute phase reactions. In a complex with MAP3K8, NFKB1/p105 represses MAP3K8-induced MAPK signaling; active MAP3K8 is released by proteasome-dependent degradation of NFKB1/p105.

#### Sequence similarities

Contains 7 ANK repeats.  
Contains 1 death domain.  
Contains 1 RHD (Rel-like) domain.

#### Domain

The C-terminus of p105 might be involved in cytoplasmic retention, inhibition of DNA-binding, and transcription activation.  
Glycine-rich region (GRR) appears to be a critical element in the generation of p50.


#### Post-translational modifications

While translation occurs, the particular unfolded structure after the GRR repeat promotes the generation of p50 making it an acceptable substrate for the proteasome. This process is known as cotranslational processing. The processed form is active and the unprocessed form acts as an inhibitor (I kappa B-like), being able to form cytosolic complexes with NF-kappa B, trapping it in the cytoplasm. Complete folding of the region downstream of the GRR repeat precludes processing.  
Phosphorylation at 'Ser-903' and 'Ser-907' primes p105 for proteolytic processing in response to TNF-alpha stimulation. Phosphorylation at 'Ser-927' and 'Ser-932' are required for BTRC/BTRCP-mediated proteolysis.  
Polyubiquitination seems to allow p105 processing.  
S-nitrosylation of Cys-61 affects DNA binding.

#### Cellular localization

Nucleus. Cytoplasm. Nuclear, but also found in the cytoplasm in an inactive form complexed to an inhibitor.

#### Images



ab114185 on a 12.5% SDS-PAGE Stained with Coomassie Blue.

SDS-PAGE - Recombinant Human NFkB p105 / p50 protein (ab114185)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

## **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors