abcam

Product datasheet

Recombinant Human RFXANK protein ab167884

1 Image

Description

Product name Recombinant Human RFXANK protein

Purity > 85 % SDS-PAGE.

ab167884 is purified using conventional chromatography techniques.

Expression system Escherichia coli

Accession <u>O14593-2</u>

Protein length Full length protein

Animal free No

Nature Recombinant

Species Human

Sequence MGSSHHHHHH SSGLVPRGSH MGSMELTQPA

EDLIQTQQTP ASELGDPEDP GEEAADGSDT VVLSLFPCTP EPVNPEPDAS VSSPQGSSLK HSTTLTNRQR GNEVSALPAT LDCDNLVNKP DERGFTPLIW ASAFGEIETV RFLLEWGADP

HILAKERESA LSLASTGGYT DIVGLLLERD VDINIYDWNG

GTPLLYAVRG NHVKCVEALL ARGADLTTEA DSGYTPMDLA VALGYRKVQQ VIENHILKLF

QSNLVPADPE

Predicted molecular weight 28 kDa including tags

Amino acids 1 to 237

Tags His tag N-Terminus

Specifications

Our Abpromise guarantee covers the use of ab167884 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications Mass Spectrometry

SDS-PAGE

Mass spectrometry MALDI-TOF

Form Liquid

Preparation and Storage

Stability and Storage

Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.

pH: 8.00

Constituents: 0.32% Tris HCI, 20% Glycerol (glycerin, glycerine), 0.88% Sodium chloride

General Info

Function Activates transcription from class II MHC promoters. Activation requires the activity of the MHC

class II transactivator (MHC2TA). May regulate other genes in the cell. RFX binds the X1 box of MHC-II promoters. Isoform RFX-B-delta5 is not involved in the positive regulation of MHC class II

genes.

Tissue specificity

Ubiquitous.

Involvement in disease

Defects in RFXANK are a cause of bare lymphocyte syndrome type 2 (BLS2) [MIM:209920]; also known as hereditary MHC class II deficiency or HLA class II-deficient combined immunodeficiency. BLS2 is a severe combined immunodeficiency disease with early onset. It is characterized by a profound defect in constitutive and interferon-gamma induced MHC II expression, absence of cellular and humoral T-cell response to antigen challenge,

hypogammaglobulinemia and impaired antibody production. The consequence include extreme

susceptibility to viral, bacterial and fungal infections.

Sequence similarities

Contains 5 ANK repeats.

Domain

The third ankyrin repeat is required for association with the two other RFX subunits; RFX5 and


RFXAP.

Cellular localization

Nucleus.

Images

(ab167884)



SDS-PAGE of ab167884 (3 μ g) under reducing condition and visualized by coomassie blue stain.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors