abcam

Product datasheet

Recombinant Human SIRP alpha protein (His tag) (Biotin) ab198765

1 Image

Description

Product name Recombinant Human SIRP alpha protein (His tag) (Biotin)

Purity >= 90 % SDS-PAGE.

Expression system HEK 293 cells

Accession P78324

Protein length Protein fragment

Animal free No

Nature Recombinant

Species Human

Sequence GVAGEEELQVIQPDKSVLVAAGETATLRCTATSLIPVGPIQ

WFRGAGPGR

ELIYNQKEGHFPRVTTVSDLTKRNNMDFSIRIGNITPADAGT

YYCVKFRK

GSPDDVEFKSGAGTELSVRAKPSAPVVSGPAARATPQH

TVSFTCESHGFS

PRDITLKWFKNGNELSDFQTNVDPVGESVSYSIHSTAKVV

LTREDVHSQV

 ${\tt ICEVAHVTLQGDPLRGTANLSETIRVPPTLEVTQQPVRAEN}$

QVNVTCQVR

KFYPQRLQLTWLENGNVSRTETASTVTENKDGTYNWMSW

LLVNVSAHRDD

VKLTCQVEHDGQPAVSKSHDLKVSAHPKEQGSNTAAEN

TGSNERGGGLND IFEAQKIEWHEHHHHHH

Predicted molecular weight 40 kDa including tags

Amino acids 27 to 370

Tags His tag C-Terminus

Additional sequence information C-terminal Avi and His tags. Enzymatically biotinylated using Avi-tag™ technology. NP_542970.

(aa 27-370 corresponds to the extracellular portion of SIRP alpha).

Conjugation Biotin

Specifications

1

Specifications

Our **Abpromise guarantee** covers the use of **ab198765** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications SDS-PAGE

Form Liquid

Preparation and Storage

Stability and Storage Shipped on Dry Ice. Store at -80°C. Avoid freeze / thaw cycle.

pH: 7.4

Constituents: 0.63% Tris HCI, 0.64% Sodium chloride, 0.02% Potassium chloride, 20% Glycerol

(glycerin, glycerine)

General Info

Function Immunoglobulin-like cell surface receptor for CD47. Acts as docking protein and induces

translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. Supports adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. May play a key role in intracellular signaling during synaptogenesis and in synaptic function (By similarity). Involved in the negative regulation of receptor tyrosine kinase-coupled cellular responses induced by cell adhesion, growth factors or insulin. Mediates negative regulation of phagocytosis, mast cell activation and dendritic cell activation. CD47 binding prevents maturation

of immature dendritic cells and inhibits cytokine production by mature dendritic cells.

Tissue specificity Ubiquitous. Highly expressed in brain. Detected on myeloid cells, but not T-cells. Detected at

lower levels in heart, placenta, lung, testis, ovary, colon, liver, small intestine, prostate, spleen,

kidney, skeletal muscle and pancreas.

Sequence similaritiesContains 2 lg-like C1-type (immunoglobulin-like) domains.

Contains 1 lg-like V-type (immunoglobulin-like) domain.

Post-translational

modifications

N-glycosylated.

and PDGF. Dephosphorylated by PTPN11.

Cellular localization Membrane.

Images



4-20% SDS-PAGE analysis of 2.7 μg ab198765 with Coomassie staining.

Note: SIRP alpha is heavily glycosylated, resulting in higher molecular weight.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors