abcam

Product datasheet

Recombinant human SMURF 2 protein ab95930

2 Images

Description

Product name Recombinant human SMURF 2 protein

Biological activity Specific Activity: 0.00049 pmol/min/μg Assay conditions: 50 μl reaction mix (4 mM ATP, 5mM

MgCl2, 1mM DTT, E1 (0.5 μ g), E2 (2.8 μ g), Biotin-Ubiquitin (0.5 μ g) and SMURF 2 (0-2 μ g) in 50mM Tris pH 7.4) incubate for 30min at 37°C, then add to streptavidin coated plates and incubate for 30min at 37°C. Add antibody against SMURF 2, incubate 1h. Finally, incubate 1h with the Europeum-labeled secondary antibody. Time resolved fluorescence measured at room temperature (Lambda exc=330 nm, Lambda em=620 nm). Specific Activity: 0.00049 pmol/min/ μ g Assay conditions: 50 μ l reaction mix (4 mM ATP, 5mM MgCl2, 1mM DTT, E1 (0.5 μ g), E2 (2.8 μ g), Biotin-Ubiquitin (0.5 μ g) and SMURF 2 (0-2 μ g) in 50mM Tris pH 7.4) incubate for 30min at 37°C, then add to streptavidin coated plates and incubate for 30min at 37°C. Add antibody against SMURF 2, incubate 1h. Finally, incubate 1h with the Europeum-labeled secondary antibody. Time resolved fluorescence measured at room temperature (Lambda exc=330 nm,

Lambda em=620 nm).

Purity > 80 % SDS-PAGE.

Affinity purified.

Expression system Baculovirus infected insect cells

Accession Q9HAU4

Protein length Protein fragment

Animal free No

Nature Recombinant

Species Human

Predicted molecular weight 72 kDa including tags

Amino acids 130 to 748

Tags DDDDK tag N-Terminus

Specifications

Our Abpromise quarantee covers the use of ab95930 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications SDS-PAGE

Functional Studies

Form Liquid

Preparation and Storage

Stability and Storage Shipped on Dry Ice. Store at -80°C.

pH: 8.00

Constituents: 0.0462% (R*,R*)-1,4-Dimercaptobutan-2,3-diol, 0.395% Tris HCl, 0.05% Tween,

20% Glycerol (glycerin, glycerine), 0.58% Sodium chloride

102 µg/ml FLAG peptide

This product is an active protein and may elicit a biological response in vivo, handle with caution.

General Info

Function E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the

form of a thioester and then directly transfers the ubiquitin to targeted substrates. Interacts with SMAD1 and SMAD7 in order to trigger their ubiquitination and proteasome-dependent degradation. In addition, interaction with SMAD7 activates autocatalytic degradation, which is prevented by interaction with SCYE1. Forms a stable complex with the TGF-beta receptor-mediated phosphorylated SMAD2 and SMAD3. In this way, SMAD2 may recruit substrates, such as SNON, for ubiquitin-mediated degradation. Enhances the inhibitory activity of SMAD7 and reduces the transcriptional activity of SMAD2. Coexpression of SMURF2 with SMAD1 results in considerable decrease in steady-state level of SMAD1 protein and a smaller decrease of

SMAD2 level.

Tissue specificity Widely expressed.

Pathway Protein modification; protein ubiquitination.

Sequence similarities Contains 1 C2 domain.

Contains 1 HECT (E6AP-type E3 ubiquitin-protein ligase) domain.

Contains 3 WW domains.

Domain The second and third WW domains are responsible for interaction with the PY-motif of R-SMAD

(SMAD1, SMAD2 and SMAD3).

The C2 domain is involved in autoinhibition of the catalytic activity by interacting with the HECT

domain.

Post-translational modifications

Auto-ubiquitinated and ubiquitinated in the presence of RNF11 and UBE2D1.

Cellular localization Nucleus. Cytoplasm. Cell membrane. Membrane raft. Cytoplasmic in the presence of SMAD7.

Co-localizes with CAV1, SMAD7 and TGF-beta receptor in membrane rafts.

Images





Image showing specific activity of ab95930.



10% SDS-PAGE

Lane 1: ab95930 (4µg)

Lane 2: Molecular weight markers

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

. Guarantee only valid for products bought direct from Abcam or one of our authorized distributors