

Product datasheet

Recombinant Human TAF15 protein ab174418

1 Image

Description

Product name	Recombinant Human TAF15 protein
Purity	> 90 % SDS-PAGE. ab174418 is purified using conventional chromatography techniques.
Expression system	Escherichia coli
Accession	<u>Q92804</u>
Protein length	Protein fragment
Animal free	No
Nature	Recombinant
Species	Human
Sequence	MGSSHHHHHH SSGLVPRGSH MGSSYHSQRE NYSHHTQDDR RDVSRYGEDN RGYGGSQGGG RGRGGYDKDG RGPMTGSSGG DRGGFKNFGG HRDYGPRTDA DSESDNSDNN TIFVQQLGEG VSTDQVGEFF KQIGIIKTNK KTGKPMINLY TDKDTGKPKG EATVSFDDPP SAKAAIDWFD GKEFHGNIK VSFATRPEF MRGGGSGGGR RGRGGYRGRG GFQGRGGDPK SGDWWCPNPS CGNMNFARRN SCNQCNEPRP EDSRPSGGDF RGRGYGGERG YR
Predicted molecular weight	30 kDa
Amino acids	148 to 406
Tags	His tag N-Terminus
Additional sequence information	NCBI Accession No. NP_631961.

Specifications

Our **Abpromise guarantee** covers the use of **ab174418** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	SDS-PAGE Mass Spectrometry
Mass spectrometry	MALDI-TOF
Form	Liquid

Preparation and Storage

Stability and Storage

Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.

pH: 8.00

Constituents: 0.32% Tris HCl, 30% Glycerol (glycerin, glycerine), 0.88% Sodium chloride, 0.02% DTT

General Info

Function

RNA and ssDNA-binding protein that may play specific roles during transcription initiation at distinct promoters. Can enter the preinitiation complex together with the RNA polymerase II (Pol II).

Tissue specificity

Ubiquitous. Observed in all fetal and adult tissues.

Involvement in disease

Note=A chromosomal aberration involving TAF15/TAF2N is found in a form of extraskeletal myxoid chondrosarcomas (EMC). Translocation t(9;17)(q22;q11) with NR4A3.

Sequence similarities

Belongs to the RRM TET family.

Contains 1 RanBP2-type zinc finger.

Contains 1 RRM (RNA recognition motif) domain.

Post-translational modifications


Dimethylated by PRMT1 at Arg-206 to asymmetric dimethylarginine. The methylation may favor nuclear localization and positive regulation of TAF15 transcriptional activity.

Phosphorylated upon DNA damage, probably by ATM or ATR.

Cellular localization

Nucleus. Cytoplasm. Shuttles from the nucleus to the cytoplasm.

Images



15% SDS-PAGE analysis of ab174418 (3µg).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors