

Product datasheet

Recombinant mouse CTLA4 protein (Active) (Biotin) ab271481

[2 Images](#)

Description

Product name	Recombinant mouse CTLA4 protein (Active) (Biotin)
Biological activity	Recombinant Mouse CTLA4 protein (Active) (Biotin): B7-1 TR-FRET IC ₅₀ = 1.8 nm.
Purity	>= 90 % SDS-PAGE.
Expression system	HEK 293 cells
Accession	<u>P09793</u>
Protein length	Protein fragment
Animal free	No
Nature	Recombinant
Species	Mouse
Sequence	EAIQVTQPSVVLASSHGVASFPCEYSPSHNTDEVRVTVLR QTNDQMTEVC ATTFTEKNTVGFLDYPFCSTGFNFSRVNLTIQGLRAVDTGL YLCKVELMYPPPYFVGMGNGTQIYVIDPEPCPDS
Predicted molecular weight	42 kDa
Amino acids	36 to 161
Tags	Avi tag C-Terminus , Fc tag C-Terminus
Additional sequence information	Extracellular domain fused to the Fc portion of human IgG1.
Conjugation	Biotin

Specifications

Our **Abpromise guarantee** covers the use of **ab271481** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	Functional Studies SDS-PAGE
Form	Liquid
Additional notes	Enzymatically biotin-labeled using Avi-tag™ technology

Preparation and Storage

Stability and Storage

Shipped on Dry Ice. Store at -80°C. Avoid freeze / thaw cycle. Store In the Dark.

pH: 7.40

Constituents: 0.13% Sodium phosphate, 0.64% Sodium chloride, 0.02% Potassium chloride, 20% Glycerol (glycerin, glycerine)

This product is an active protein and may elicit a biological response in vivo, handle with caution.

General Info

Function

Inhibitory receptor acting as a major negative regulator of T-cell responses. The affinity of CTLA4 for its natural B7 family ligands, CD80 and CD86, is considerably stronger than the affinity of their cognate stimulatory coreceptor CD28.

Tissue specificity

Widely expressed with highest levels in lymphoid tissues. Detected in activated T-cells where expression levels are 30- to 50-fold less than CD28, the stimulatory coreceptor, on the cell surface following activation.

Involvement in disease

Genetic variation in CTLA4 influences susceptibility to systemic lupus erythematosus (SLE) [MIM:152700]. SLE is a chronic, inflammatory and often febrile multisystemic disorder of connective tissue. It affects principally the skin, joints, kidneys and serosal membranes. SLE is thought to represent a failure of the regulatory mechanisms of the autoimmune system. Note=Genetic variations in CTLA4 may influence susceptibility to Graves disease, an autoimmune disorder associated with overactivity of the thyroid gland and hyperthyroidism. Genetic variation in CTLA4 is the cause of susceptibility to diabetes mellitus insulin-dependent type 12 (IDDM12) [MIM:601388]. A multifactorial disorder of glucose homeostasis that is characterized by susceptibility to ketoacidosis in the absence of insulin therapy. Clinical features are polydipsia, polyphagia and polyuria which result from hyperglycemia-induced osmotic diuresis and secondary thirst. These derangements result in long-term complications that affect the eyes, kidneys, nerves, and blood vessels. Genetic variation in CTLA4 is the cause of susceptibility to celiac disease type 3 (CELIAC3) [MIM:609755]. It is a multifactorial disorder of the small intestine that is influenced by both environmental and genetic factors. It is characterized by malabsorption resulting from inflammatory injury to the mucosa of the small intestine after the ingestion of wheat gluten or related rye and barley proteins. In its classic form, celiac disease is characterized in children by malabsorption and failure to thrive.

Sequence similarities

Contains 1 Ig-like V-type (immunoglobulin-like) domain.


Post-translational modifications

N-glycosylation is important for dimerization. Phosphorylation at Tyr-201 prevents binding to the AP-2 adapter complex, blocks endocytosis, and leads to retention of CTLA4 on the cell surface.

Cellular localization


Cell membrane. Exists primarily as an intracellular antigen whose surface expression is tightly regulated by restricted trafficking to the cell surface and rapid internalisation and.

Images



Recombinant Mouse CTLA4 protein (Active) (Biotin): B7-1 TR-FRET $IC_{50} = 1.8 \text{ nM}$.

Functional Studies - Recombinant mouse CTLA4 protein (Active) (Biotin) (ab271481)



SDS-PAGE analysis of 2 µg ab271481.

This protein runs at a higher MW by SDS-PAGE due to glycosylation.

SDS-PAGE - Recombinant mouse CTLA4 protein (Active) (Biotin) (ab271481)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Promise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors