# abcam

# Product datasheet

# Streptavidin (HRP) ab7403

\*\*\* \* \* \* 6 Abreviews 150 References 2 Images

### **Description**

Product name Streptavidin (HRP)

**Biological activity** Binds to Biotin. Dissociation constant has not been measured.

Purity > 95 % SDS-PAGE.

Chromatographically pure, a single band by SDS-PAGE. Streptavidin-HRP was prepared from chromatographically purified streptavidin. Streptavidin Peroxidase conjugate was assayed by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase and anti-

Streptavidin.

Expression system Native

Accession P22629

Protein length Full length protein

Animal free No

**Nature** Native

**Species** Streptomyces avidinii

Conjugation HRP

# **Specifications**

Our Abpromise guarantee covers the use of ab7403 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

**Applications** Dot blot

**ELISA** 

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemistry (Frozen sections)

Immunomicroscopy
Immunocytochemistry

Western blot

Form Liquid

Additional notes Horseradish peroxidase is conjugated to the streptavidin tetramer at ~1:1 molar ratio.

This product has been assayed against 1.0  $\mu$ g of Biotinylated lgG in a standard capture ELISA using a **peroxidase substrate** as **ABTS** ab142041 as a substrate for 30 minutes at room

1

temperature. A working dilution of 1:15,000 to 1:60,000 of the reconstitution concentration is suggested for this product. Optimal titers for other applications should be determined by the researcher.

# **Preparation and Storage**

#### Stability and Storage

Shipped at 4°C. Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.

Preservative: 0.01% Gentamicin sulphate

Constituents: 0.424% Potassium phosphate, 0.87% Sodium chloride, BSA

This product is an active protein and may elicit a biological response in vivo, handle with caution.

#### **General Info**


#### Relevance

Streptavidin is a tetrameric protein purified from Streptomyces sp. that binds very tightly to the vitamin biotin with a Kd of  $\sim$  10-14 mol/l. The high affinity recognition of biotin and biotinylated molecules has made streptavidin one of the most important components in diagnostics and laboratory kits.

#### **Cellular localization**

Cytoplasmic

#### **Images**




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Streptavidin (HRP) (ab7403)

IHC image of Histone H1 staining in a section of formalin-fixed paraffin-embedded [human normal colon]\*. The section was pretreated using pressure cooker heat mediated antigen retrieval with sodium citrate buffer (pH6) for 30mins. The section was then incubated with <a href="mailto:ab11080">ab11080</a>, 1/1000 dilution, for 15 mins at room temperature. A goat anti-mouse biotinylated secondary antibody (<a href="mailto:ab6788">ab6788</a>, 1/1000 dilution), was used to detect the primary, and visualized using an HRP conjugated ABC system. Streptavidin HRP was used, ab7403 at a 1/10000 dilution. DAB was used as the chromogen (<a href="mailto:ab103723">ab103723</a>), diluted 1/100 and incubated for 10min at room temperature. The section was then counterstained with haematoxylin and mounted with DPX. The inset negative control image is taken from an identical assay without primary antibody.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

\*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre



The wells were coated with <u>ab200699</u> at 1µg/ml at 50µl/well overnight at 4°C, followed by a 5% BSA blocking step for 2h RT. Human lgG3 (<u>ab138703</u>) was then added starting at 20 µg/ml and plasma/serum at 1:500 and gradually diluted 1:4, 50µl/well for 2h. <u>Ab201248</u> was then added at 1:10,000 dilution, 50µl/well for 2h. A HRP-streptavidin (ab7403) was used at 1:10,000 dilution for 1h.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.com/abpromise">https://www.abcam.com/abpromise</a> or contact our technical team.

#### Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors