

## Product datasheet

# Biotinylated Goat anti-Mouse IgG (H+L) (Ready to Use) ab64255

[23 References](#)   [1 Image](#)

### Overview

<b>Product name</b>	Biotinylated Goat anti-Mouse IgG (H+L) (Ready to Use)
<b>Tested applications</b>	<b>Suitable for:</b> Sandwich ELISA, IHC-P
<b>General notes</b>	<p><b>STAINING PROTOCOL:</b></p> <p>Deparaffinize and rehydrate formalin-fixed paraffin-embedded tissue section.</p> <p>Add enough drops of an appropriate Hydrogen Peroxide Block to cover the sections.</p> <p>Incubate for 10 minutes. Wash 2 times in buffer.</p> <p>Perform appropriate pretreatment if required. Wash 3 times in buffer.</p> <p>Apply an appropriate Protein Block and incubate for 5 minutes at room temperature to block nonspecific background staining. Wash 1 time in buffer.</p> <p>Apply primary antibody and incubate according to manufacturer's protocol.</p> <p>Wash 4 times in buffer. Apply Biotinylated Goat anti Mouse IgG (H+L) solution and incubate for 10 minutes at room temperature. Wash 4 times in buffer.</p> <p>Apply enzyme labeled streptavidin and incubate according to manufacturer's protocol.</p> <p>Rinse 4 times in buffer. Place slide in appropriate chromogenic substrate and incubate until desired reaction is achieved. Rinse 4 times in buffer.</p> <p>Add enough drops of Hematoxylin to cover the section. Incubate for 1 minute.</p> <p>Rinse 7-8 times in tap water. Add Mounting Medium to cover the section.</p>

This reagent is intended to be used in a labeled streptavidin-biotin immunoenzymatic antigen detection system. This technique involves the sequential incubation of the specimen with an unconjugated primary antibody specific to the target antigen, a biotinylated secondary antibody which reacts with the primary antibody, enzyme-labeled streptavidin, and substrate-chromogen.

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Store at +4°C.
<b>Storage buffer</b>	<p>pH: 7.2</p> <p>Preservative: 0.0985% Sodium azide</p> <p>Constituent: PBS</p>
<b>Reagent notes</b>	This reagent is intended to be used in a labeled streptavidin-biotin immunoenzymatic antigen detection system. This technique involves the sequential incubation of the specimen with an

unconjugated primary antibody specific to the target antigen, a biotinylated secondary antibody which reacts with the primary antibody, enzyme-labeled streptavidin, and substrate-chromogen.

<b>Cellular localization</b>	Secreted
<b>Cellular localization</b>	Secreted


## Applications

**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab64255 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
<b>Sandwich ELISA</b>		Use at an assay dependent concentration. PubMed: 23754944
<b>IHC-P</b>		Use at an assay dependent dilution.

## Images



Human esophagus tissue section stained with biotinylated Goat anti-Mouse IgG (H+L) (ab64255).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Biotinylated Goat anti-Mouse IgG (H+L) (Ready to Use) (ab64255)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors