# abcam

### Product datasheet

## Goat Anti-Rabbit IgG Fc (Alexa Fluor® 555) ab150090

3 References 3 Images

Overview

Product name Goat Anti-Rabbit IgG Fc (Alexa Fluor® 555)

Host species Goat
Target species Rabbit

**Tested applications** Suitable for: ICC/IF, Flow Cyt, ELISA, IHC-P, IHC-Fr

**Immunogen** Other Immunogen Type corresponding to Rabbit IgG.

**Conjugation** Alexa Fluor® 555. Ex: 555nm, Em: 565nm

**Properties** 

Form Liquid

**Storage instructions** Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Stable for 12 months at -20°C. Store In the Dark.

Storage buffer Preservative: 0.02% Sodium azide

Constituents: 23% Glycerol (glycerin, glycerine), PBS, 1% BSA

Purity Immunogen affinity purified

**Purification notes**This antibody was isolated by affinity chromatography using antigen coupled to agarose beads.

**Clonality** Polyclonal

**Isotype** IgG

General notes Alexa Fluor<sup>®</sup> is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific

Company. The Alexa Fluor<sup>®</sup> dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor<sup>®</sup> dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor<sup>®</sup> dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact

Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or

1

#### outlicensing@thermofisher.com.

#### **Applications**


The Abpromise guarantee

Our <u>Abpromise guarantee</u> covers the use of ab150090 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


| Application | Abreviews | Notes                                    |
|-------------|-----------|--|
| ICC/IF      |           | 1/200 - 1/1000.                          |
| Flow Cyt    |           | 1/2000.                                  |
| ELISA       |           | Use at an assay dependent concentration. |
| IHC-P       |           | Use at an assay dependent concentration. |
| IHC-Fr      |           | Use at an assay dependent concentration. |

#### **Images**




Immunocytochemistry/ Immunofluorescence - Goat
Anti-Rabbit IgG Fc (Alexa Fluor® 555) (ab150090)


ICC/IF image of <u>ab6046</u> stained HeLa cells. The cells were 100% methanol fixed (5 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (<u>ab6046</u>, 1 $\mu$ g/ml) overnight at +4°C. The secondary antibody (yellow) was ab150090 Alexa Fluor® 555 goat anti-rabbit lgG Fc used at 2 $\mu$ g/ml for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43 $\mu$ M.



Immunocytochemistry/ Immunofluorescence - Goat Anti-Rabbit IgG Fc (Alexa Fluor® 555) (ab150090)



HeLa cells showing negative staining by ICC/IF using only secondary antibody. The cells were 100% methanol fixed (5 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The secondary antibody (yellow) was ab150090 Alexa Fluor® 555 goat anti-rabbit IgG Fc used at 2µg/ml for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

• Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.com/abpromise">https://www.abcam.com/abpromise</a> or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors