

Product datasheet

Mouse lung tissue lysate - total protein (14 days) ab7268

1 References

Overview

Product name	Mouse lung tissue lysate - total protein (14 days)
General notes	Source: Outbred NIH-Swiss Albino mice (14 days old)

The mouse lung tissue was collected from fourteen (14) day-old mice. The lysate was prepared by homogenization in modified RIPA buffer (50 mM Tris-HCl, pH 7.4, 1% Triton X-100, 0.2% sodium deoxycholate, 0.2% sodium dodecylsulfate (SDS), 1 mM sodium ethylenediaminetetraacetate, 1 mM phenylmethylsulfonyl flouride, 5 µg/ml of aprotinin, 5 µg/ml of leupeptin). Tissue and cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The lysate was boiled for 5 min in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% SDS, 0.01% bromophenol blue) containing 5% b-mercaptoethanol.

Abcam has not and does not intend to apply for the REACH Authorisation of customers' uses of products that contain European Authorisation list (Annex XIV) substances.
It is the responsibility of our customers to check the necessity of application of REACH Authorisation, and any other relevant authorisations, for their intended uses.

Tested applications	Suitable for: WB
----------------------------	-------------------------

Properties

Mycoplasma free	Yes
Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.2 Constituents: 60.05% Water, 12.5% Glycerol (glycerin, glycerine), 9% Tris HCl, 7.7% DTT, 4.4% Sodium chloride, 1% Triton-X-100, 1% Sodium deoxycholate, 0.15% EDTA disodium salt, 0.5% Aprotinin, 0.5% Leupeptin hemisulfate, 0.09% PMSF, 0.01% Bromophenol blue

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab7268 in the following tested applications.
The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent dilution. Mouse lung tissue lysate is ready to load on SDS-PAGE for Western blotting. It is recommended to load 10 µg to 20 µg per lane for mini gel

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors