

Product datasheet

Protein A Sepharose® ab193256

[7 References](#) [2 Images](#)

Overview

Product name	Protein A Sepharose®
Sample type	Cell culture supernatant, Serum, Cell culture media, Ascites Fluid
Product overview	Contents: Supplied as a 50% slurry in 20% Ethanol.

Features:

High binding capacity = Binding of IgG ≥ 16 mg human or rabbit IgG/mL Protein A-Sepharose®.

Minimal leaching of the ligand

Flow Rate Tested* = 2.07 mL/min.

*Test condition: = Calculations based on the time required to pass 18 mL of water through 2 mL settled beads (column diameter 1.5 cm).

Usage = Reusable for up to 10 times without significant loss of binding capacity.

Store beads at 4°C.

The beads may be damaged above 40°C.

DO NOT FREEZE.

Wash beads 3 times with 3x bead volume of desired buffer before use.

Applications:

- Purification of monoclonal and polyclonal antibodies from culture media, serum, ascites fluid or hybridoma supernatants.

- Isolation of antibody/antigen complexes in immunoprecipitation experiments, since only the Fc region is involved in antibody binding and the Fab region is available for binding antigen.

Sepharose is a registered trademark of GE Healthcare

Notes

This product is manufactured by BioVision, an Abcam company and was previously called 6501 Protein A Sepharose. 6501-5 is the same size as the 5 ml size of ab193256.

Protein A Sepharose[®] beads are prepared by covalently coupling recombinant Protein A to 6% cross-linked Sepharose[®] beads. Protein A is a genetically engineered protein containing five IgG-binding regions of native Protein A. The cell wall binding region, albumin binding region and other non-specific regions have been eliminated from the recombinant Protein A to ensure maximum specific IgG binding. The coupling technique is optimized to give a higher binding capacity for IgG and minimum leaching of recombinant Protein A. The IgG binding capacity of Protein A Sepharose[®] is ≥ 16 mg human or rabbit IgG per mL of wet beads. Protein A Sepharose[®] beads display high chemical and physical stability as well as high flow rate, hydrophilicity and high gel strength. This product can be used for IgG purification and immunoprecipitation.

Tested applications

Suitable for: IP, Purification

Properties

Storage instructions

Store at +4°C. Please refer to protocols.

Components	1 ml	5 ml	25 ml	100 ml
Protein A Sepharose [®]	1 x 1ml	1 x 5ml	1 x 25ml	1 x 100ml

Applications


The Abpromise guarantee

Our [Abpromise guarantee](#) covers the use of ab193256 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IP		Use at an assay dependent concentration. Isolation of antibody/antigen complexes in immunoprecipitation experiments, since only the Fc region is involved in antibody binding and the Fab region is available for binding antigen.
Purification		Use at an assay dependent concentration. Purification of monoclonal and polyclonal antibodies from culture media, serum, ascites fluid or hybridoma supernatants.


Images



Immunoprecipitation - Protein A Sepharose®
(ab193256)

Image from Arnold et al., J Exp Clin Cancer Res., 39(1):205; doi: 10.1186/s13046-020-01712-w. Reproduced under the Creative Commons license <https://creativecommons.org/licenses/by/4.0/>

Overexpression of TTLL4 in MDA-MB231 cells increases MT-glutamylation. Polyglutamylated proteins were immunoprecipitated from control and TTLL4 plus cells, using an antibody against polyglutamylation modification (GT335). Cell lysates ("input"), supernatants ("S/N") from cell lysates incubated with GT335-coupled beads and proteins bound to GT335 coupled beads ("beads") were analyzed by Western blotting for β -tubulin and NAP-1 levels. IgG signals served as loading control. The lower band appearing in the NAP-1 blot is a residue signal of β -tubulin antibody because the same membrane was used to probe against all 3 proteins.



Immunoprecipitation - Protein A Sepharose®
(ab193256)

Image from O'Leary et al., Sci Rep., 8(1):15360; doi: 10.1038/s41598-018-33840-x. Reproduced under the Creative Commons license <https://creativecommons.org/licenses/by/4.0/>

Immunoprecipitation of resistin from obese subcutaneous adipose conditioned medium secretome (OB ACM) improves myogenesis. Resistin was immunoprecipitated from OB ACM using resistin antibody-agarose bead conjugates (OB ACM - resistin IP). IgG isotype antibody control-agarose bead conjugates were used on the same samples as a control (OB ACM). Resistin protein is detected by immunoblotting of resistin-antibody lysates but not IgG control lysates following immunoprecipitation.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors