

Product datasheet

Anti-RanGAP1 antibody [EPR3295] - BSA and Azide free ab239907

Recombinant RabMAb

10 Images

Overview

Product name	Anti-RanGAP1 antibody [EPR3295] - BSA and Azide free
Description	Rabbit monoclonal [EPR3295] to RanGAP1 - BSA and Azide free
Host species	Rabbit
Tested applications	Suitable for: Flow Cyt, ICC/IF, IP, WB, IHC-P
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Synthetic peptide within Human RanGAP1 aa 1-100. The exact sequence is proprietary. Database link: P46060
Positive control	IHC-P: Human breast carcinoma, Human testis, Human bladder carcinoma, Mouse testis and Rat liver tissue. ICC/IF: mouse hepatocyte, and MCF7 cells. FC: Jurkat cells
General notes	<p>ab239907 is the carrier-free version of ab92360. This format is designed for use in antibody labeling, including fluorochromes, metal isotopes, oligonucleotides, enzymes.</p> <p>Our carrier-free formats are supplied in a buffer free of BSA, sodium azide and glycerol for higher conjugation efficiency.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>ab239907 is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm.</p> <p><i>Maxpar® is a trademark of Fluidigm Canada Inc.</i></p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.</p> <p>We are constantly working hard to ensure we provide our customers with best in class antibodies. As a result of this work we are pleased to now offer this antibody in purified format. We are in the process of updating our datasheets. The purified format is</p>

designated 'PUR' on our product labels. If you have any questions regarding this update, please contact our Scientific Support team.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR3295
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab239907** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		Use at an assay dependent concentration. ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ICC/IF		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration. Predicted molecular weight: 64 kDa.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol.

Target

Function	GTPase activator for the nuclear Ras-related regulatory protein Ran, converting it to the putatively inactive GDP-bound state.
Tissue specificity	Highly expressed in brain, thymus and testis.
Sequence similarities	Belongs to the RNA1 family. Contains 6 LRR (leucine-rich) repeats.
Post-translational modifications	Phosphorylated occurs before nuclear envelope breakdown and continues throughout mitosis. Phosphorylated by the M-phase kinase cyclin B/Cdk1, in vitro. Differential timing of dephosphorylation occurs during phases of mitosis. The phosphorylated form remains associated with RANBP2/NUP358 and the SUMO E2-conjugating enzyme, UBC9, on nuclear pore complex


(NPC) disassembly and during mitosis.

Sumoylated with SUMO1. Sumoylation is necessary for targeting to the nuclear envelope (NE), and for association with mitotic spindles and kinetochores during mitosis. Also required for interaction with RANBP2 and is mediated by UBC9.

Cellular localization

Cytoplasm. Nucleus membrane. Chromosome, centromere, kinetochore. Cytoplasm, cytoskeleton, spindle pole. Cytoplasmic during interphase. Targeted to the nuclear rim after sumoylation. During mitosis, associates with mitotic spindles. Association with kinetochores appears soon after nuclear envelope breakdown and persists until late anaphase. Mitotic location also requires sumoylation.


Images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Rat liver tissue sections labeling RanGAP1 with purified [ab92360](#) at 1/500 dilution (0.22 µg/ml). Perform heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab92360](#)).


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RanGAP1 antibody [EPR3295] - BSA and Azide free ([ab239907](#))



Immunocytochemistry/ Immunofluorescence analysis of MCF7 (Human breast adenocarcinoma epithelial cell) cells labeling RanGAP1 with purified [ab92360](#) at 1/100 dilution (1.1 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with [ab195889](#) Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) at 1/200 (2.5 µg/ml) dilution. Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) was used as the secondary antibody at 1/1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.

Immunocytochemistry/ Immunofluorescence - Anti-RanGAP1 antibody [EPR3295] - BSA and Azide free ([ab239907](#))

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab92360](#)).



Immunoprecipitation - Anti-RanGAP1 antibody
[EPR3295] - BSA and Azide free (ab239907)

[ab92360](#) (purified) at 1/20 dilution (0.5ug) immunoprecipitating RanGAP1 in HeLa whole cell lysates.

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates 10ug


Lane 2 (+): [ab92360](#) & HeLa whole cell lysates

Lane 3 (-): Rabbit monoclonal IgG ([ab172730](#)) instead of [ab92360](#) in HeLa whole cell lysates

For western blotting, VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) was used at 1/1000 dilution.

Blocking and diluting buffer: 5% NFD/MTBST.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab92360](#)).



Flow Cytometry - Anti-RanGAP1 antibody
[EPR3295] - BSA and Azide free (ab239907)

Flow Cytometry analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labeling RanGAP1 with purified [ab92360](#) at 1/20 dilution (10 µg/ml) (red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab92360](#)).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RanGAP1 antibody [EPR3295] - BSA and Azide free (ab239907)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Mouse testis tissue sections labeling RanGAP1 with purified [ab92360](#) at 1/500 dilution (0.22 µg/ml). Perform heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab92360](#)).



Immunocytochemistry/ Immunofluorescence - Anti-RanGAP1 antibody [EPR3295] - BSA and Azide free (ab239907)

Immunofluorescence staining of MCF7 cells with purified [ab92360](#) at a working dilution of 1/500, counter-stained with DAPI. The secondary antibody was an Alexa Fluor[®] 488 conjugated goat anti-rabbit ([ab150077](#)), used at a dilution of 1/1000. The cells were fixed in 4% PFA and permeabilized using 0.1% Triton X 100. The negative control is shown in bottom right hand panel - for the negative control, PBS was used instead of the primary antibody.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab92360](#)).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RanGAP1 antibody [EPR3295] - BSA and Azide free (ab239907)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human bladder carcinoma tissue sections labeling RanGAP1 with purified [ab92360](#) at 1/500 dilution (0.22 µg/ml). Perform heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab92360](#)).




Immunocytochemistry/ Immunofluorescence - Anti-RanGAP1 antibody [EPR3295] - BSA and Azide free (ab239907)

This image is courtesy of an anonymous Abreview.

[ab92360](#) staining RanGAP1 in mouse hepatocyte cells by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with paraformaldehyde, permeabilized, and blocked with 2% BSA for 2 hours at 22°C. Samples were incubated with primary antibody (1/100 in blocking buffer) for 18 hours at 4°C. An Alexa Fluor® 488-conjugated goat anti-rabbit IgG polyclonal (1/10000) was used as the secondary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab92360](#)).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RanGAP1 antibody [EPR3295] - BSA and Azide free (ab239907)

[ab92360](#) at 1/100 dilution staining RanGAP1 in paraffin-embedded (1) Human breast carcinoma tissue and (2) Human testis tissue by immunohistochemistry.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab92360](#)).

Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol.



Flow Cytometry - Anti-RanGAP1 antibody
[EPR3295] - BSA and Azide free (ab239907)

Overlay histogram showing Jurkat cells stained with [ab92360](#) (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody ([ab92360](#), 1/100 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) ([ab96899](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line). Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in Jurkat cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab92360](#)).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors