

Product datasheet

Recombinant Human ApoER2 protein ab152817

1 Image

Description

| | | |
|-----------------------------------|---|--|
| Product name | Recombinant Human ApoER2 protein | |
| Expression system | Wheat germ | |
| Accession | Q14114 | |
| Protein length | Protein fragment | |
| Animal free | No | |
| Nature | Recombinant | |
| Species | Human | |
| Sequence | KKTCADSDFTCDNGHCIHERWKCDGEEECPDGSESEA TCTKQVCPAEKL SCGPTSHKCVPASWRCDGEKDCEGGADEAGCATLCAP H | |
| Predicted molecular weight | 35 kDa including tags | |
| Amino acids | 83 to 170 | |

Specifications

Our [Abpromise guarantee](#) covers the use of **ab152817** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| | |
|---------------------|--------------|
| Applications | SDS-PAGE |
| | Western blot |
| | ELISA |
| Form | Liquid |

Additional notes


Preparation and Storage

| | |
|------------------------------|--|
| Stability and Storage | Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles. pH: 8.00 Constituents: 0.31% Glutathione, 0.79% Tris HCl |
|------------------------------|--|

General Info

| | |
|---|---|
| Function | Cell surface receptor for Reelin (RELN) and apolipoprotein E (apoE)-containing ligands. LRP8 participates in transmitting the extracellular Reelin signal to intracellular signaling processes, by binding to DAB1 on its cytoplasmic tail. Reelin acts via both the VLDL receptor (VLDLR) and LRP8 to regulate DAB1 tyrosine phosphorylation and microtubule function in neurons. LRP8 has higher affinity for Reelin than VLDLR. LRP8 is thus a key component of the Reelin pathway which governs neuronal layering of the forebrain during embryonic brain development. Binds the endoplasmic reticulum resident receptor-associated protein (RAP). Binds dimers of beta 2-glycoprotein I and may be involved in the suppression of platelet aggregation in the vasculature. Highly expressed in the initial segment of the epididymis, where it affects the functional expression of clusterin and phospholipid hydroperoxide glutathione peroxidase (PHGPx), two proteins required for sperm maturation. May also function as an endocytic receptor. |
| Tissue specificity | Expressed mainly in brain and placenta. Also expressed in platelets and megakaryocytic cells. Not expressed in the liver. |
| Involvement in disease | Defects in LRP8 are a cause of myocardial infarction type 1 (MCI1) [MIM:608446]. A condition defined by the irreversible necrosis of heart muscle secondary to prolonged ischemia. |
| Sequence similarities | Belongs to the LDLR family. Contains 2 EGF-like domains. Contains 7 LDL-receptor class A domains. Contains 5 LDL-receptor class B repeats. |
| Domain | The cytoplasmic domain is involved in the binding of DAB1 and in the recruitment of JNK-interacting proteins. Isoforms, which lack part of the cytoplasmic domain, are unable to recruit members of the family of JNK interacting proteins (JIP) to the cytoplasmic tail. |
| Post-translational modifications | O-glycosylated. Some alternatively spliced isoforms lack the O-linked sugar domain. Undergoes sequential, furin and gamma-secretase dependent, proteolytic processing, resulting in the extracellular release of the entire ligand-binding domain as a soluble polypeptide and in the intracellular domain (ICD) release into the cytoplasm. The gamma-secretase-dependent proteolytical processing occurs after the bulk of the extracellular domain has been shed, in a furin-dependent manner, in alternatively spliced isoforms carrying the furin cleavage site. Hypoglycosylation (mainly hypo-O-glycosylation) leads to increased extracellular cleavage, which in turn results in accelerating release of the intracellular domain (ICD) by the gamma-secretase. The resulting receptor fragment is able to inhibit Reelin signaling and in particular the Reelin-induced DAB1 phosphorylation. Tyrosine phosphorylated upon apoE binding. Ubiquitinated by MYLIP leading to degradation. |
| Cellular localization | Cell membrane. Secreted. Isoforms that contain the exon coding for a furin-type cleavage site are proteolytically processed, leading to a secreted receptor fragment. |

Images



12.5% SDS-PAGE analysis of ab152817 stained with Coomassie Blue.

SDS-PAGE - Recombinant Human ApoER2 protein
(ab152817)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors