

Product datasheet

Recombinant human CD200 / OX2 protein (Fc Chimera) ab214617

Description

Product name	Recombinant human CD200 / OX2 protein (Fc Chimera)	
Biological activity	Shows the biological function of the CD200 / OX2 moiety and exerts a prolonged circulating half-life caused by the modified Fc domain.	
Purity	≥ 98 % SDS-PAGE.	
Endotoxin level	< 0.060 Eu/μg	
Expression system	CHO cells	
Accession	P41217	
Protein length	Protein fragment	
Animal free	No	
Nature	Recombinant	
Species	Human	
Sequence	QVQVVTQDEREQLYTPASLKCSLQNAQEALIVTWQKKKA VSPENMVFSE NHGVVIQPAYKDKINITQLGLQNSTITFWNITLEDEGCMCL FNTFGFGK ISGTACLTVYVQPIVSLHYKFSEDHLNITCSATARPAPMVF WKVPRSGIE NSTVTLSHPNGTTSVTSILHIKDPKNQVGKEVICQVLHLGTV TDFKQTVN KGYWFS	
Predicted molecular weight	23 kDa	
Amino acids	31 to 236	
Additional sequence information	Extracellular domain fused to the N-terminus of the Fc region of a mutant Human IgG1 (NP_005935.4).	

Specifications

Our [Abpromise guarantee](#) covers the use of **ab214617** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	SDS-PAGE
	Functional Studies

Form	Lyophilized
Additional notes	Non-lytic: Acts as a long lasting fusion protein which only binds to the receptor. Mutations to the complement (C1q) and FcγR I binding sites of the IgGs Fc fragment render the fusion proteins incapable of antibody directed cytotoxicity (ADCC) and complement directed cytotoxicity (CDC).

Preparation and Storage

Stability and Storage	Shipped at 4°C. Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle. Constituent: 100% PBS Lyophilized from 0.2µm filtered solution. This product is an active protein and may elicit a biological response in vivo, handle with caution.
Reconstitution	Reconstitute at 100µg/ml in sterile PBS.

General Info

Function	Costimulates T-cell proliferation. May regulate myeloid cell activity in a variety of tissues.
Sequence similarities	Contains 1 Ig-like C2-type (immunoglobulin-like) domain. Contains 1 Ig-like V-type (immunoglobulin-like) domain.
Cellular localization	Cell membrane.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors