

Product datasheet

Recombinant human Cystatin SA/CST2 protein
ab180061

2 Images

Description

Product name	Recombinant human Cystatin SA/CST2 protein	
Biological activity	Measured by its ability to inhibit active Cathepsin L cleavage of a fluorogenic peptide substrate Z-LR-AMC. The IC ₅₀ value is < 10.0 nM.	
Purity	> 95 % SDS-PAGE.	
Endotoxin level	< 1.000 Eu/μg	
Expression system	HEK 293 cells	
Accession	P09228	
Protein length	Full length protein	
Animal free	No	
Nature	Recombinant	
Species	Human	
Sequence	WSPQEEDRIIEGGYDADLNDERVQRALHFVISEYNKATED EYRRLLRV LRAREQIVGGVNYFFDIEVGRITCKTSQPNLDTCAFHEQPE LQKKQLCSF QYEVPWEDRMSLVNSRCQEA	
Predicted molecular weight	15 kDa including tags	
Amino acids	21 to 141	
Tags	His tag C-Terminus	
Additional sequence information	(AAH62679)	

Specifications

Our [Abpromise guarantee](#) covers the use of **ab180061** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	Functional Studies
	SDS-PAGE
Form	Lyophilized

Preparation and Storage

Stability and Storage

Shipped at 4°C. Store at 4°C (stable for up to 12 months). Store at -80°C. Avoid freeze / thaw cycle. For long term storage it is recommended to add a carrier protein on reconstitution (0.1% HSA or BSA).

pH: 7.4

Constituents: 5% Trehalose, 0.3% Tris, 0.88% Sodium chloride

Lyophilized from 0.22 µm filtered solution.

This product is an active protein and may elicit a biological response in vivo, handle with caution.

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile deionized water to a final concentration of 400 µg/ml. Solubilize for 30 to 60 minutes at room temperature with occasional gentle mixing.

General Info

Function

Thiol protease inhibitor.

Tissue specificity

Expressed in submandibular and sublingual saliva but not in parotid saliva (at protein level).
Expressed in submandibular gland and parotid gland.


Sequence similarities

Belongs to the cystatin family.

Cellular localization


Secreted.

Images



Human Cystatin SA (His Tag) on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

SDS-PAGE - Recombinant human Cystatin
SA/CST2 protein (ab180061)



SDS-PAGE analysis of ab180061 in reducing conditions. Gel stained overnight with Coomassie Blue. DTT-reduced protein migrates as 15-17 kDa.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors