

Product datasheet

Recombinant human EGFR protein (Mutant) ab208473

2 Images

Description

Product name	Recombinant human EGFR protein (Mutant)
Biological activity	The specific activity of ab208473 was determined to be 90 nmol/min/mg.
Purity	> 75 % Densitometry.
Expression system	Baculovirus infected Sf9 cells
Accession	P00533
Protein length	Protein fragment
Animal free	No
Nature	Recombinant
Species	Human
Sequence	<p>SGEAPNQALLRILKETEFKKIKVLGSGAFGTVYKGLWIP EGEKVKIPVAI KEATSPKANKEILDEAYVMASVDNPHVCRLLGICLTST VQLITQLMPFGC LLDYVREHKDNIGSQYLLNWCVQIAKGMNYLEDRLVH RDLAARNVLVKT PQHVKITDFGLAKLLGAEKEKYHAEGGKVPKWMAL SILHRIYTHQSDV WSYGVTVWELMTFGSKPYDGIPASEISSILEKGERLPQ PPICTIDVYMIM VKCWMIDADSRPKFRELIIEFSKMARDPQRYLVIQGDE RMHLPSPTDSNF YRALMDEEDMDDVDADEYLIPQQGFFSSPSTSRTP LSSLSATSNNSTV ACIDRNLQSCPIKEDSFLQRYSSDPTGALTEDSIDDT FLPVPEYINQSV PKRPAGSVQNPVYHNQPLNPAPSRDPHYQDPHSTAV GNPEYLNTVQPTCV NSTFDSPAHWAKGSHQISLDNPDYQQDFFPKEAKP NGIFKGSTAENAEYLRVAPQSSEFIGA</p>
Predicted molecular weight	87 kDa including tags
Amino acids	695 to 1210
Tags	GST tag N-Terminus

Additional sequence information Mutation (d747-749) (NM_005228).

Specifications

Our [Abpromise guarantee](#) covers the use of **ab208473** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	SDS-PAGE Functional Studies
Form	Liquid

Preparation and Storage

Stability and Storage	Shipped on Dry Ice. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle. pH: 7.5 Constituents: 0.79% Tris HCl, 0.87% Sodium chloride, 0.31% Glutathione, 0.003% EDTA, 0.002% PMSF, 0.004% DTT, 25% Glycerol This product is an active protein and may elicit a biological response in vivo, handle with caution.
------------------------------	--

General Info


Function	Receptor tyrosine kinase binding ligands of the EGF family and activating several signaling cascades to convert extracellular cues into appropriate cellular responses. Known ligands include EGF, TGFA/TGF-alpha, amphiregulin, epigen/EPGN, BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding EGF. Ligand binding triggers receptor homo- and/or heterodimerization and autophosphorylation on key cytoplasmic residues. The phosphorylated receptor recruits adapter proteins like GRB2 which in turn activates complex downstream signaling cascades. Activates at least 4 major downstream signaling cascades including the RAS-RAF-MEK-ERK, PI3 kinase-AKT, PLCgamma-PKC and STATs modules. May also activate the NF-kappa-B signaling cascade. Also directly phosphorylates other proteins like RGS16, activating its GTPase activity and probably coupling the EGF receptor signaling to the G protein-coupled receptor signaling. Also phosphorylates MUC1 and increases its interaction with SRC and CTNNB1/beta-catenin. Isoform 2 may act as an antagonist of EGF action.
Tissue specificity	Ubiquitously expressed. Isoform 2 is also expressed in ovarian cancers.
Involvement in disease	Lung cancer Inflammatory skin and bowel disease, neonatal, 2
Sequence similarities	Belongs to the protein kinase superfamily. Tyr protein kinase family. EGF receptor subfamily. Contains 1 protein kinase domain.
Post-translational modifications	Phosphorylation at Ser-695 is partial and occurs only if Thr-693 is phosphorylated. Phosphorylation at Thr-678 and Thr-693 by PRKD1 inhibits EGF-induced MAPK8/JNK1 activation. Dephosphorylation by PTPRJ prevents endocytosis and stabilizes the receptor at the plasma membrane. Autophosphorylation at Tyr-1197 is stimulated by methylation at Arg-1199 and enhances interaction with PTPN6. Autophosphorylation at Tyr-1092 and/or Tyr-1110 recruits STAT3. Dephosphorylated by PTPN1 and PTPN2. Monoubiquitinated and polyubiquitinated upon EGF stimulation; which does not affect tyrosine kinase activity or signaling capacity but may play a role in lysosomal targeting. Polyubiquitin

linkage is mainly through 'Lys-63', but linkage through 'Lys-48', 'Lys-11' and 'Lys-29' also occurs. Deubiquitination by OTUD7B prevents degradation. Ubiquitinated by RNF115 and RNF126. Methylated. Methylation at Arg-1199 by PRMT5 stimulates phosphorylation at Tyr-1197.

Cellular localization


Secreted and Cell membrane. Endoplasmic reticulum membrane. Golgi apparatus membrane. Nucleus membrane. Endosome. Endosome membrane. Nucleus. In response to EGF, translocated from the cell membrane to the nucleus via Golgi and ER. Endocytosed upon activation by ligand. Colocalized with GPER1 in the nucleus of estrogen agonist-induced cancer-associated fibroblasts (CAF).

Images



SDS-PAGE analysis of ab208473.

SDS-PAGE - Recombinant human EGFR protein (Mutant) (ab208473)



Sample Kinase Activity assay plot using ab208473, showing the specific activity to be 90 nmol/min/mg.

Functional Studies - Recombinant human EGFR protein (Mutant) (ab208473)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors