

Product datasheet

Recombinant Human Frizzled 2/FZD2 protein ab191625

1 Image

Description

Product name	Recombinant Human Frizzled 2/FZD2 protein	
Purity	> 95 % SDS-PAGE.	
Endotoxin level	< 1.000 Eu/g	
Expression system	HEK 293 cells	
Accession	Q14332	
Protein length	Protein fragment	
Animal free	No	
Nature	Recombinant	
Species	Human	
Sequence	QFHGEKGISIPDHGFCQPISIPLCTDIAYNQTIMPNLLGHTNQ EDAGLEV HQFYPLVKVQCSPFLRFFLCMYAPVCTVLEQAIPPCRSI CERARQGCEA LMNKFGFQWPERLRCEHFPRHGAEQICVGQNHS	
Predicted molecular weight	42 kDa including tags	
Amino acids	24 to 156	
Additional sequence information	Fused with Fc fragment of Human IgG1 at the C terminus	

Specifications

Our [Abpromise guarantee](#) covers the use of **ab191625** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	SDS-PAGE
Form	Lyophilized
Additional notes	This product was previously labelled as Frizzled 2

Preparation and Storage

Stability and Storage

Shipped at 4°C. Store at 4°C prior to reconstitution. Store at -80°C. Avoid freeze / thaw cycle.

pH: 7.4

Constituents: 0.61% Tris, 0.75% Glycine, Sodium chloride, L-Arginine

Lyophilized from 0.22 µm filtered solution. Normally trehalose is added as protectant before lyophilization.

Reconstitution

Reconstitute with sterile deionized water to a concentration of 200 µg/ml.

General Info

Function

Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues.

Tissue specificity

Widely expressed. In the adult, mainly found in heart, placenta, skeletal muscle, lung, kidney, pancreas, prostate, testis, ovary and colon. In the fetus, expressed in brain, lung and kidney. Low levels in fetal liver.

Sequence similarities

Belongs to the G-protein coupled receptor Fz/Smo family.
Contains 1 FZ (frizzled) domain.


Domain

Lys-Thr-X-X-X-Trp motif is involved in the activation of the Wnt/beta-catenin signaling pathway. The FZ domain is involved in binding with Wnt ligands.

Cellular localization

Membrane.

Images



SDS-PAGE analysis of DTT-reduced ab191625, stained overnight with Coomassie Blue. DTT-reduced Frizzled 2/FZD2 migrates as 45-50 kDa due to glycosylation.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors