

Product datasheet

Recombinant human HAUSP / USP7 protein (Active)
ab269127

2 Images

Description

Product name	Recombinant human HAUSP / USP7 protein (Active)
Biological activity	ab269127 specific activity was determined to be 4 nmol/mon/mg in a DUB assay using recombinant human ubiquitin-based proluciferin substrate.
Purity	> 75 % SDS-PAGE.
Expression system	Escherichia coli
Accession	Q93009
Protein length	Protein fragment
Animal free	No
Nature	Recombinant
Species	Human
Molecular weight information	Approx 68 kDa by SDS-PAGE
Amino acids	25 to 580
Tags	His tag N-Terminus
Additional sequence information	NM_003470

Specifications

Our [Abpromise guarantee](#) covers the use of **ab269127** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	Functional Studies SDS-PAGE
Form	Liquid

Preparation and Storage

Stability and Storage	Shipped on Dry Ice. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle. pH: 7.00 Preservative: 1.02% Imidazole
------------------------------	--

Constituents: 0.82% Sodium phosphate, 1.74% Sodium chloride, 0.002% PMSF, 0.004% DTT, 25% Glycerol (glycerin, glycerine)

This product is an active protein and may elicit a biological response in vivo, handle with caution.

General Info

Function

Hydrolase that deubiquitinates target proteins such as FOXO4, p53/TP53, MDM2, ERCC6, DNMT1, UHRF1, PTEN and DAXX (PubMed:11923872, PubMed:15053880, PubMed:16964248, PubMed:18716620, PubMed:25283148). Together with DAXX, prevents MDM2 self-ubiquitination and enhances the E3 ligase activity of MDM2 towards p53/TP53, thereby promoting p53/TP53 ubiquitination and proteasomal degradation. Deubiquitinates p53/TP53, preventing degradation of p53/TP53, and enhances p53/TP53-dependent transcription regulation, cell growth repression and apoptosis (PubMed:25283148). Deubiquitinates p53/TP53 and MDM2 and strongly stabilizes p53/TP53 even in the presence of excess MDM2, and also induces p53/TP53-dependent cell growth repression and apoptosis. Deubiquitination of FOXO4 in presence of hydrogen peroxide is not dependent on p53/TP53 and inhibits FOXO4-induced transcriptional activity. In association with DAXX, is involved in the deubiquitination and translocation of PTEN from the nucleus to the cytoplasm, both processes that are counteracted by PML. Involved in cell proliferation during early embryonic development. Involved in transcription-coupled nucleotide excision repair (TC-NER) in response to UV damage: recruited to DNA damage sites following interaction with KIAA1530/UVSSA and promotes deubiquitination of ERCC6, preventing UV-induced degradation of ERCC6. Contributes to the overall stabilization and trans-activation capability of the herpesvirus 1 trans-acting transcriptional protein ICP0/VMW110 during HSV-1 infection. Involved in maintenance of DNA methylation via its interaction with UHRF1 and DNMT1: acts by mediating deubiquitination of UHRF1 and DNMT1, preventing their degradation and promoting DNA methylation by DNMT1 (PubMed:21745816). Exhibits a preference towards 'Lys-48'-linked ubiquitin chains. Increases regulatory T-cells (Treg) suppressive capacity by deubiquitinating and stabilizing the transcription factor FOXP3 which is crucial for Treg cell function (PubMed:23973222).

Tissue specificity

Widely expressed. Overexpressed in prostate cancer.

Sequence similarities

Belongs to the peptidase C19 family.
Contains 1 MATH domain.
Contains 1 USP domain.

Domain

The C-terminus plays a role in its oligomerization.


Post-translational modifications

Isoform 1: Phosphorylated. Isoform 1 is phosphorylated at positions Ser-18 and Ser-963. Isoform 2: Not phosphorylated.
Isoform 1: Polyneddylated. Isoform 2: Not Polyneddylated.
Isoform 1 and isoform 2: Not sumoylated.
Isoform 1 and isoform 2: Polyubiquitinated by herpesvirus 1 trans-acting transcriptional protein ICP0/VMW110; leading to its subsequent proteasomal degradation. Isoform 1: Ubiquitinated at Lys-869.

Cellular localization


Nucleus. Cytoplasm. Nucleus, PML body. Present in a minority of ND10 nuclear bodies. Association with ICP0/VMW110 at early times of infection leads to an increased proportion of USP7-containing ND10. Colocalizes with ATXN1 in the nucleus. Colocalized with DAXX in speckled structures. Colocalized with PML and PTEN in promyelocytic leukemia protein (PML) nuclear bodies.

Images



Functional Studies - Recombinant human HAUSP /
USP7 protein (Tagged) (ab269127)

ab269127 specific activity was determined to be 4 nmol/mon/mg in a DUB assay using recombinant human ubiquitin-based proluciferin substrate.



SDS-PAGE - Recombinant human HAUSP / USP7
protein (Tagged) (ab269127)

SDS-PAGE analysis of ab269127.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors