

Product datasheet

Recombinant human HDAC3 protein ab101663

5 Images

Description

Product name	Recombinant human HDAC3 protein
Biological activity	Specific activity 125 RLU/min/ng.
Purity	> 70 % SDS-PAGE. ab101663 is >70% pure as assessed by densitometry.
Expression system	Baculovirus
Accession	O15379
Protein length	Full length protein
Animal free	No
Nature	Recombinant
Species	Human
Predicted molecular weight	83 kDa including tags
Amino acids	1 to 428

Specifications

Our [Abpromise guarantee](#) covers the use of **ab101663** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	SDS-PAGE Functional Studies
Form	Liquid

Preparation and Storage

Stability and Storage	Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles. pH: 7.50 Constituents: 0.307% Glutathione, 0.00174% PMSF, 0.00385% DTT, 0.79% Tris HCl, 0.00292% EDTA, 25% Glycerol (glycerin, glycerine), 0.87% Sodium chloride This product is an active protein and may elicit a biological response in vivo, handle with caution.
------------------------------	--

General Info

Function

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Probably participates in the regulation of transcription through its binding to the zinc-finger transcription factor YY1; increases YY1 repression activity. Required to repress transcription of the POU1F1 transcription factor. Acts as a molecular chaperone for shuttling phosphorylated NR2C1 to PML bodies for sumoylation.

Tissue specificity

Widely expressed.

Sequence similarities

Belongs to the histone deacetylase family, HD type 1 subfamily.


Post-translational modifications

Sumoylated in vitro.

Cellular localization

Nucleus.

Images




The specific activity of HDAC3 (ab101663) was determined to be 144 RLU/min/mg as per activity assay protocol

Functional Studies - Recombinant human HDAC3 protein (ab101663)




SDS PAGE analysis of ab101663

SDS-PAGE - Recombinant human HDAC3 protein (ab101663)




SDS PAGE analysis of ab101663

SDS-PAGE - Recombinant human HDAC3 protein (ab101663)



The specific activity of ab101663 was determined to be 125 RLU/min/ng.

Functional Studies - Recombinant human HDAC3 protein (ab101663)



The purity of HDAC3 was determined to be >70% as assessed by densitometry.

The approximate molecular weight is ~83 kDa.

SDS-PAGE - Recombinant human HDAC3 protein (ab101663)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors