

Product datasheet

# Recombinant human ZAP70 protein ab60963

2 Images

## Description

---

<b>Product name</b>	Recombinant human ZAP70 protein
<b>Purity</b>	> 95 % Densitometry. Affinity purified.
<b>Expression system</b>	Baculovirus infected Sf9 cells
<b>Accession</b>	<b><u>P43403</u></b>
<b>Protein length</b>	Full length protein
<b>Animal free</b>	No
<b>Nature</b>	Recombinant
<b>Species</b>	Human
<b>Predicted molecular weight</b>	96 kDa including tags
<b>Amino acids</b>	1 to 619
<b>Tags</b>	GST tag N-Terminus

## Specifications

---

Our **Abpromise guarantee** covers the use of **ab60963** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>Applications</b>	Functional Studies
<b>Form</b>	Liquid
<b>Additional notes</b>	<b><u>ab204877</u></b> (Poly (4:1 Glu, Tyr) peptide) can be utilized as a substrate for assessing kinase activity

## Preparation and Storage

---


<b>Stability and Storage</b>	Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles. pH: 7.50 Constituents: 0.0038% EGTA, 0.00174% PMSF, 0.00385% DTT, 0.79% Tris HCl, 0.00292% EDTA, 25% Glycerol, 0.87% Sodium chloride  This product is an active protein and may elicit a biological response in vivo, handle with caution.
------------------------------	--

## General Info


## General info

<b>Function</b>	Plays a role in T-cell development and lymphocyte activation. Essential for TCR-mediated IL-2 production. Isoform 1 induces TCR-mediated signal transduction, isoform 2 does not.
<b>Tissue specificity</b>	Expressed in T- and natural killer cells.
<b>Involvement in disease</b>	Defects in ZAP70 are the cause of selective T-cell defect (STD) [MIM:176947]. STD is an autosomal recessive form of severe combined immunodeficiency characterized by a selective absence of CD8-type T-cells.
<b>Sequence similarities</b>	Belongs to the protein kinase superfamily. Tyr protein kinase family. SYK/ZAP-70 subfamily. Contains 1 protein kinase domain. Contains 2 SH2 domains.
<b>Domain</b>	The SH2 domains bind to the phosphorylated tyrosine-based activation motif (TAM) of CD3Z and the non-canonical phosphorylated tyrosine-based activation motif (TAM) of RHOH.
<b>Post-translational modifications</b>	Phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation. Tyr-319 phosphorylation is essential for full activity.
<b>Cellular localization</b>	Cytoplasm. Cell membrane. After antigen stimulation, isoform 1 concentrates at the immunological synapse and isoform 2 remains cytoplasmic. Co-localizes together with RHOH in the immunological synapse. RHOH is required for its proper localization to the cell membrane and cytoskeleton fractions in the thymocytes.

## Images



SDS PAGE analysis of ab60963



The specific activity of ZAP70 (ab60963) was determined to be 129 nmol/min/mg as per activity assay protocol

Functional Studies - Recombinant human ZAP70 protein (ab60963)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors