

Product datasheet

Anti-S100A9 antibody [EPR3555] ab92507

Recombinant RabMAb

15 References 7 Images

Overview

Product name	Anti-S100A9 antibody [EPR3555]
Description	Rabbit monoclonal [EPR3555] to S100A9
Host species	Rabbit
Tested applications	Suitable for: WB, IHC-P, ICC/IF, Flow Cyt Unsuitable for: IP
Species reactivity	Reacts with: Human
Immunogen	Synthetic peptide within Human S100A9 aa 1-100. The exact sequence is proprietary.
Positive control	Breast cancer, Human tonsil, Human spleen and peripheral blood lymphocytes lysates, Human breast carcinoma tissue and Human spleen tissue.IF: DU145 cell lineFlow Cyt: Jurkat cells
General notes	Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information [see here](#).

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).

Reproducibility is key to advancing scientific discovery and accelerating scientists' next breakthrough.

Abcam is leading the way with our range of recombinant antibodies, knockout-validated antibodies and knockout cell lines, all of which support improved reproducibility.

We are also planning to innovate the way in which we present recommended applications and species on our product datasheets, so that only applications & species that have been tested in our own labs, our suppliers or by selected trusted collaborators are covered by our Abpromise[™] guarantee.

In preparation for this, we have started to update the applications & species that this product is Abpromise guaranteed for.

We are also updating the applications & species that this product has been “predicted to work with,” however this information is not covered by our Abpromise guarantee.

Applications & species from publications and Abreviews that have not been tested in our own labs or in those of our suppliers are not covered by the Abpromise guarantee.

Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, as well as customer reviews and Q&As.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
Storage buffer	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 9% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA, 50% Tissue culture supernatant
Purity	Tissue culture supernatant
Clonality	Monoclonal
Clone number	EPR3555
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab92507** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/1000 - 1/5000. Predicted molecular weight: 13 kDa.
IHC-P		1/250 - 1/500. Perform heat mediated antigen retrieval via the microwave method before commencing with IHC staining protocol. Boil slides seven times for five minutes each in pH 6 citrate buffer.
ICC/IF		1/100 - 1/250.
Flow Cyt		1/200.

Application notes	Is unsuitable for IP.
--------------------------	-----------------------

Target

Function	Calcium-binding protein. Has antimicrobial activity towards bacteria and fungi. Important for resistance to invasion by pathogenic bacteria. Up-regulates transcription of genes that are under the control of NF-kappa-B. Plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS) (By similarity). Promotes tubulin polymerization when
-----------------	---

unphosphorylated. Promotes phagocyte migration and infiltration of granulocytes at sites of wounding. Plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1. Extracellular calprotectin binds to target cells and promotes apoptosis. Antimicrobial and proapoptotic activity is inhibited by zinc ions.

Tissue specificity

Expressed by macrophages in acutely inflamed tissues and in chronic inflammation. Detected in peripheral blood leukocytes, in neutrophils and granulocytes. Detected at sites of vascular inflammation (at protein level). Also expressed in epithelial cells constitutively or induced during dermatoses.

Sequence similarities

Belongs to the S-100 family.
Contains 2 EF-hand domains.


Post-translational modifications

Phosphorylated. Phosphorylation inhibits activation of tubulin polymerization.

Cellular localization

Secreted. Cytoplasm. Cytoplasm > cytoskeleton. Cell membrane. Associates with tubulin filaments in activated monocytes. Targeted to the cell surface upon calcium influx. Released from blood leukocytes upon exposure to CSF2/GM-CSF, bacterial lipopolysaccharide (LPS) and during inflammatory processes. Serum levels are high in patients suffering from chronic inflammation.


Images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-S100A9 antibody [EPR3555] (ab92507)


Anti-S100A9 antibody [EPR3555] (ab92507)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human tonsil tissue labelling S100A9 with ab92507 at a dilution of 1:500. Heat mediated antigen retrieval was performed using AR9 antigen retrieval solution, and microwave treatment for 15 min at 20% power. Anti-Rabbit/Mouse HRP polymer (PerkinElmer Opal Polymer HRP Ms Plus Rb) was used as secondary antibody. Opal tyramide amplification was performed using Opal 520 fluorophore. Counterstained with DAPI stain. Image scanned with Vectra 3.0 and analyzed via Phenochart software. This image was courteously provided by Dr. Houssein Abdul Sater, Georgia Cancer Center.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-S100A9 antibody [EPR3555] (ab92507)

ab92507, at a 1/250 dilution, staining S100A9 in paraffin embedded Human breast carcinoma tissue by Immunohistochemistry.



Western blot - Anti-S100A9 antibody [EPR3555] (ab92507)

All lanes : Anti-S100A9 antibody [EPR3555] (ab92507) at 1/1000 dilution

Lane 1 : Breast cancer lysate

Lane 2 : Human tonsil lysate

Lane 3 : Human spleen lysate

Lane 4 : Peripheral blood lymphocytes lysate

Lysates/proteins at 10 µg per lane.


Secondary

All lanes : HRP labelled goat anti-rabbit IgG at 1/2000 dilution

Predicted band size: 13 kDa


Observed band size: 14 kDa

[why is the actual band size different from the predicted?](#)




Flow Cytometry - Anti-S100A9 antibody [EPR3555] (ab92507)

Flow cytometry analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labeling S100A9 (red) with purified ab92507 at a 1/200 dilution (10ug/mL). Cells were fixed with 80% methanol and permeabilized with 0.1% Tween-20. A goat anti rabbit IgG (Alexa Fluor® 488) ([ab150077](#)) was used as the secondary antibody at a 1/2000 dilution. Black - Rabbit monoclonal IgG (Black) ([ab172730](#)). Blue (unlabeled control) - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-S100A9 antibody [EPR3555] (ab92507)

ab92507, at a 1/250 dilution, staining S100A9 in paraffin embedded Human spleen tissue by Immunohistochemistry.



ICC/IF image of ab92507 stained DU145 cells. The cells were 4% paraformaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab92507, 1/200) overnight at +4°C. The secondary antibody (green) was ab96899, DyLight® 488 goat anti-rabbit IgM (H+L) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

Immunocytochemistry/ Immunofluorescence - Anti-S100A9 antibody [EPR3555] (ab92507)

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-S100A9 antibody [EPR3555] (ab92507)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors