

Product datasheet

Anti-Serine racemase antibody [EPR16682] ab200833

Recombinant RabMAb

3 Images

Overview

Product name	Anti-Serine racemase antibody [EPR16682]
Description	Rabbit monoclonal [EPR16682] to Serine racemase
Host species	Rabbit
Tested applications	Suitable for: WB, IP Unsuitable for: ICC/IF
Species reactivity	Reacts with: Human
Immunogen	Synthetic peptide within Human Serine racemase aa 50-150. The exact sequence is proprietary. Database link: Q9GZT4
Positive control	WB: LNCaP, U-87 MG and HEK293 whole cell lysates; Human fetal brain lysate. IP: HEK293 and LNCaP whole cell lysates.

General notes

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMab[®] patents](#).

This product is a [recombinant rabbit monoclonal antibody](#).

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.05% BSA
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR16682
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab200833** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/2000. Detects a band of approximately 37 kDa (predicted molecular weight: 37 kDa).
IP		1/50.
Application notes		Is unsuitable for ICC/IF.

Target

Function

Catalyzes the synthesis of D-serine from L-serine. D-serine is a key coagonist with glutamate at NMDA receptors. Has dehydratase activity towards both L-serine and D-serine.

Tissue specificity

Brain: expressed at high levels in hippocampus and corpus callosum, intermediate levels in substantia nigra and caudate, and low levels in amygdala, thalamus, and subthalamic nuclei. Expressed in heart, skeletal muscle, kidney and liver.


Sequence similarities

Belongs to the serine/threonine dehydratase family.

Post-translational modifications

S-nitrosylated, leading to decrease the enzyme activity.

Images



All lanes : Anti-Serine racemase antibody [EPR16682] (ab200833) at 1/5000 dilution

Lane 1 : LNCaP (Human prostate cancer cell line) whole cell lysate

Lane 2 : Human fetal brain lysate

Lane 3 : U-87 MG (Human glioblastoma-astrocytoma epithelial cell line) whole cell lysate

Lane 4 : HEK293 (Human embryonic kidney) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated

Predicted band size: 37 kDa

Observed band size: 37 kDa

Exposure time: 3 minutes

Blocking/Dilution Buffer: 5% NFDm/TBST.




Immunoprecipitation - Anti-Serine racemase antibody [EPR16682] (ab200833)

Serine racemase was immunoprecipitated from 1mg of HEK293 (Human embryonic kidney) whole cell lysate with ab200833 at 1/50 dilution. Western blot was performed from the immunoprecipitate using ab200833 at 1/1000 dilution. Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG, was used as secondary antibody at 1/1500 dilution.

Lane 1: HEK293 whole cell lysate 10 μ g (Input). Lane 2: ab200833 IP in HEK293 whole cell lysate. Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab200833 in HEK293 whole cell lysate. Blocking and dilution buffer and concentration: 5% NFDm/TBST.

Exposure time: 30 seconds



Immunoprecipitation - Anti-Serine racemase antibody [EPR16682] (ab200833)

Serine racemase was immunoprecipitated from 1mg of LNCaP (Human prostate cancer cell line) whole cell lysate with ab200833 at 1/50 dilution. Western blot was performed from the immunoprecipitate using ab200833 at 1/1000 dilution. Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG, was used as secondary antibody at 1/1500 dilution.

Lane 1: LNCaP whole cell lysate 10 μ g (Input). Lane 2: ab200833 IP in LNCaP whole cell lysate. Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab200833 in LNCaP whole cell lysate. Blocking and dilution buffer and concentration: 5% NFDm/TBST.

Exposure time: 10 seconds

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors