

Product datasheet

Anti-Slow Skeletal Myosin Heavy chain antibody - N-terminal ab228353

1 Image

Overview

Product name	Anti-Slow Skeletal Myosin Heavy chain antibody - N-terminal
Description	Rabbit polyclonal to Slow Skeletal Myosin Heavy chain - N-terminal
Host species	Rabbit
Tested applications	Suitable for: WB
Species reactivity	Reacts with: Mouse Predicted to work with: Human 
Immunogen	Synthetic peptide within Human Slow Skeletal Myosin Heavy chain (N terminal). The exact sequence is proprietary. Conjugated to a protein carrier. Database link: P12883
Positive control	WB: Mouse muscle tissue extract.
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.00 Preservative: 0.01% Thimerosal (merthiolate) Constituents: 1.21% Tris, 0.75% Glycine, 10% Glycerol (glycerin, glycerine)
Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

Applications


The Abpromise guarantee Our [Abpromise guarantee](#) covers the use of ab228353 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/500 - 1/3000. Predicted molecular weight: 223 kDa.

Target

Function	Muscle contraction.
Tissue specificity	Both wild type and variant Gln-403 are detected in skeletal muscle (at protein level).
Involvement in disease	<p>Defects in MYH7 are the cause of cardiomyopathy familial hypertrophic type 1 (CMH1) [MIM:192600]. Familial hypertrophic cardiomyopathy is a hereditary heart disorder characterized by ventricular hypertrophy, which is usually asymmetric and often involves the interventricular septum. The symptoms include dyspnea, syncope, collapse, palpitations, and chest pain. They can be readily provoked by exercise. The disorder has inter- and intrafamilial variability ranging from benign to malignant forms with high risk of cardiac failure and sudden cardiac death.</p> <p>Defects in MYH7 are the cause of myopathy myosin storage (MYOMS) [MIM:608358]. In this disorder, muscle biopsy shows type 1 fiber predominance and increased interstitial fat and connective tissue. Inclusion bodies consisting of the beta cardiac myosin heavy chain are present in the majority of type 1 fibers, but not in type 2 fibers.</p> <p>Defects in MYH7 are the cause of scapuloperoneal myopathy MYH7-related (SPMM) [MIM:181430]; also known as scapuloperoneal syndrome myopathic type. SPMM is a progressive muscular atrophy beginning in the lower legs and affecting the shoulder region earlier and more severely than distal arm.</p> <p>Defects in MYH7 are a cause of cardiomyopathy dilated type 1S (CMD1S) [MIM:613426]. Dilated cardiomyopathy is a disorder characterized by ventricular dilation and impaired systolic function, resulting in congestive heart failure and arrhythmia. Patients are at risk of premature death.</p> <p>Defects in MYH7 are the cause of myopathy distal type 1 (MPD1) [MIM:160500]. MPD1 is a muscular disorder characterized by early-onset selective weakness of the great toe and ankle dorsiflexors, followed by weakness of the finger extensors. Mild proximal weakness occasionally develops years later after the onset of the disease.</p>
Sequence similarities	<p>Contains 1 IQ domain.</p> <p>Contains 1 myosin head-like domain.</p>
Domain	<p>The rodlike tail sequence is highly repetitive, showing cycles of a 28-residue repeat pattern composed of 4 heptapeptides, characteristic for alpha-helical coiled coils.</p> <p>Each myosin heavy chain can be split into 1 light meromyosin (LMM) and 1 heavy meromyosin (HMM). It can later be split further into 2 globular subfragments (S1) and 1 rod-shaped subfragment (S2).</p>
Cellular localization	Cytoplasm > myofibril. Thick filaments of the myofibrils.

Images



Anti-Slow Skeletal Myosin Heavy chain antibody - N-terminal (ab228353) at 1/2000 dilution + Mouse muscle tissue extract at 50 μ g

Developed using the ECL technique.

Predicted band size: 223 kDa

5% SDS-PAGE

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors