

Product datasheet

Anti-TBR2 / Eomes antibody [EPR19012] - BSA and Azide free ab222226

Recombinant RabMAb

[7 Images](#)

Overview

Product name	Anti-TBR2 / Eomes antibody [EPR19012] - BSA and Azide free
Description	Rabbit monoclonal [EPR19012] to TBR2 / Eomes - BSA and Azide free
Host species	Rabbit
Tested applications	Suitable for: IHC-Fr, IHC-P, ICC
Species reactivity	Reacts with: Mouse Does not react with: Human
Immunogen	Recombinant fragment aa 250-500. The exact sequence is proprietary. Database link: O54839
Positive control	IHC-P: Mouse hippocampus and E14 cortex tissues; IHC-Fr: Mouse E14.5 cortex and hippocampus tissues; ICC: Mouse primary neuron cells.
General notes	<p>Ab222226 is the carrier-free version of ab183991. This format is designed for use in antibody labeling, including fluorochromes, metal isotopes, oligonucleotides, enzymes.</p> <p>Our carrier-free formats are supplied in a buffer free of BSA, sodium azide and glycerol for higher conjugation efficiency.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>ab222226 is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm.</p> <p><i>Maxpar® is a trademark of Fluidigm Canada Inc.</i></p> <p>Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.</p> <p>Reproducibility is key to advancing scientific discovery and accelerating scientists' next breakthrough.</p> <p>Abcam is leading the way with our range of recombinant antibodies, knockout-validated antibodies and knockout cell lines, all of which support improved reproducibility.</p> <p>We are also planning to innovate the way in which we present recommended applications and species on our product datasheets, so that only applications & species that have been tested in</p>

our own labs, our suppliers or by selected trusted collaborators are covered by our Abpromise™ guarantee.

In preparation for this, we have started to update the applications & species that this product is Abpromise guaranteed for.

We are also updating the applications & species that this product has been “predicted to work with,” however this information is not covered by our Abpromise guarantee.

Applications & species from publications and Abreviews that have not been tested in our own labs or in those of our suppliers are not covered by the Abpromise guarantee.

Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, as well as customer reviews and Q&As.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze.
Storage buffer	pH: 7.2 Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR19012
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab222226** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-Fr		Use at an assay dependent concentration. Antigen retrieval: Heated citrate solution (10mM citrate PH 6.0 + 0.05% Tween-20).
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ICC		Use at an assay dependent concentration.

Target

Function	Functions as a transcriptional activator playing a crucial role during development. Functions in trophoblast differentiation and later in gastrulation, regulating both mesoderm delamination and endoderm specification. Plays a role in brain development being required for the specification
-----------------	--

and the proliferation of the intermediate progenitor cells and their progeny in the cerebral cortex. Also involved in the differentiation of CD8+ T-cells during immune response regulating the expression of lytic effector genes.

Tissue specificity

Expressed in CD8+ T-cells.

Involvement in disease

Note=A translocation t(3;10)(p24;q23) located 215 kb 3' to the EOMES gene but leading to loss of its expression was identified in a large consanguineous family. Homozygous silencing produces microcephaly associated with corpus callosum agenesis, bilateral polymicrogyria, ventricular dilatation and a small cerebellum.

Sequence similarities

Contains 1 T-box DNA-binding domain.


Developmental stage

Detected at 7 weeks of development in the forebrain floorplate of the CNS. Expressed within the mantle layer and migrating neuroblasts of the telencephalon at 12.5 weeks of development.

Cellular localization

Nucleus.

Images



Immunohistochemical analysis of paraffin-embedded mouse hippocampus tissue labeling TBR2 / Eomes with [ab183991](#) at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.


Nucleus staining on intermediate neuronal progenitors of mouse hippocampus is observed [PMID: 22553033].

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is [ab97051](#) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab183991](#)).


Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunocytochemistry - Anti-TBR2 / Eomes antibody [EPR19012] - BSA and Azide free (ab222226)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized mouse primary neuron cells labelling TBR2 / Eomes with [ab183991](#) at 1/500 dilution, followed by [ab150077](#) Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (2 µg/mL) (Green). Confocal image showing nuclear staining in mouse primary neuron cell. Confocal scanning Z step was set as 0.3 µm followed by image processing with maximum Z projection. [ab11267](#) Anti-MAP2 mouse monoclonal antibody was used to counterstain tubulin at 1/500 dilution (4 µg/mL) followed by [ab150120](#) Goat Anti-Mouse IgG H&L (Alexa Fluor® 594) at 1/1000 dilution (2 µg/mL) (Red). The nuclear counterstain was DAPI (Blue).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab183991](#)).



Immunohistochemistry (Frozen sections) - Anti-TBR2 / Eomes antibody [EPR19012] - BSA and Azide free (ab222226)


Immunohistochemical analysis of 4% paraformaldehyde-fixed, 0.2% Triton X-100 permeabilized frozen mouse E14.5 cortex tissue labeling TBR2 / Eomes with [ab183991](#) at 1/1000 dilution, followed by Goat Anti-Rabbit IgG (Alexa Fluor® 488) ([ab150077](#)) secondary antibody at 1/1000 dilution (green).

Specific staining on E14.5 mouse cortex is observed.

The nuclear counterstain is DAPI (blue).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is [ab150077](#) at 1/1000 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab183991](#)).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TBR2 / Eomes antibody [EPR19012] - BSA and Azide free (ab222226)

Immunohistochemical analysis of paraffin-embedded mouse E14 cortex tissue labeling TBR2 / Eomes with [ab183991](#) at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.


Nucleus staining on the cortex of mouse E14 is observed [PMID: 22553033].

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is [ab97051](#) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab183991](#)).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Frozen sections) - Anti-TBR2 / Eomes antibody [EPR19012] - BSA and Azide free (ab222226)


Immunohistochemical analysis of 4% paraformaldehyde-fixed, 0.2% Triton X-100 permeabilized frozen mouse hippocampus tissue labeling TBR2 / Eomes with [ab183991](#) at 1/1000 dilution, followed by Goat Anti-Rabbit IgG (Alexa Fluor® 488) ([ab150077](#)) secondary antibody at 1/1000 dilution (green).

Specific staining on adult mouse hippocampus is observed.

The nuclear counterstain is DAPI (blue).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is [ab150077](#) at 1/1000 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab183991](#)).



Immunohistochemical analysis of 4% paraformaldehyde-fixed, 0.2% Triton X-100 permeabilized frozen adult mouse kidney tissue labeling TBR2 / Eomes with [ab183991](#) at 1/1000 dilution, followed by Goat Anti-Rabbit IgG (Alexa Fluor® 488) ([ab150077](#)) secondary antibody at 1/1000 dilution (green).

Negative control tissue; very weak staining is due to autofluorescence of the kidney section.

The nuclear counterstain is DAPI (blue).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is [ab150077](#) at 1/1000 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab183991](#)).

Immunohistochemistry (Frozen sections) - Anti-TBR2 / Eomes antibody [EPR19012] - BSA and Azide free ([ab222226](#))

Why choose a recombinant antibody?

 <p>Research with confidence Consistent and reproducible results</p>	 <p>Long-term and scalable supply Recombinant technology</p>
 <p>Success from the first experiment Confirmed specificity</p>	 <p>Ethical standards compliant Animal-free production</p>

Anti-TBR2 / Eomes antibody [EPR19012] - BSA and Azide free ([ab222226](#))

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors