

## Product datasheet

# Anti-TNF alpha antibody [EPR20972] (Alexa Fluor® 647) ab237354

Recombinant RabMAb

[2 Images](#)

### Overview

<b>Product name</b>	Anti-TNF alpha antibody [EPR20972] (Alexa Fluor® 647)
<b>Description</b>	Rabbit monoclonal [EPR20972] to TNF alpha (Alexa Fluor® 647)
<b>Host species</b>	Rabbit
<b>Conjugation</b>	Alexa Fluor® 647. Ex: 652nm, Em: 668nm
<b>Tested applications</b>	<b>Suitable for:</b> ICC/IF, Flow Cyt
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse
<b>Immunogen</b>	Recombinant fragment within Mouse TNF alpha aa 1 to the C-terminus. The exact sequence is proprietary. Database link: <a href="#">P06804</a>
<b>Positive control</b>	ICC/IF: LPS (100 ng/ml, 7hours) and BFA (1ug/ml, 3hours) treated RAW264.7 cells. Flow Cyt: LPS stimulated RAW264.7 cells.
<b>General notes</b>	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb® patents</a>.</p> <p>Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research.</p>

For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or [outlicensing@thermofisher.com](mailto:outlicensing@thermofisher.com).

## Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle. Store In the Dark.
<b>Storage buffer</b>	pH: 7.4 Preservative: 0.02% Sodium azide Constituents: 30% Glycerol, 1% BSA, PBS
<b>Purity</b>	Protein A purified
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	EPR20972
<b>Isotype</b>	IgG

## Applications

Our [Abpromise guarantee](#) covers the use of **ab237354** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		1/100. This product gave a positive signal in LPS (100 ng/ml, 7hours) and BFA (1ug/ml, 3hours) treated Raw264.7 fixed with 4% formaldehyde (10 min).
Flow Cyt		1/500.

## Target

<b>Function</b>	Cytokine that binds to TNFRSF1A/TNFR1 and TNFRSF1B/TNFR2. It is mainly secreted by macrophages and can induce cell death of certain tumor cell lines. It is potent pyrogen causing fever by direct action or by stimulation of interleukin-1 secretion and is implicated in the induction of cachexia, Under certain conditions it can stimulate cell proliferation and induce cell differentiation.
<b>Involvement in disease</b>	Genetic variations in TNF are a cause of susceptibility psoriatic arthritis (PSORAS) [MIM:607507]. PSORAS is an inflammatory, seronegative arthritis associated with psoriasis. It is a heterogeneous disorder ranging from a mild, non-destructive disease to a severe, progressive, erosive arthropathy. Five types of psoriatic arthritis have been defined: asymmetrical oligoarthritis characterized by primary involvement of the small joints of the fingers or toes; asymmetrical arthritis which involves the joints of the extremities; symmetrical polyarthritis characterized by a rheumatoidlike pattern that can involve hands, wrists, ankles, and feet; arthritis mutilans, which is a rare but deforming and destructive condition; arthritis of the sacroiliac joints and spine (psoriatic spondylitis).

## Sequence similarities

Belongs to the tumor necrosis factor family.

## Post-translational modifications

The soluble form derives from the membrane form by proteolytic processing.

The membrane form, but not the soluble form, is phosphorylated on serine residues.


Dephosphorylation of the membrane form occurs by binding to soluble TNFRSF1A/TNFR1.

O-glycosylated; glycans contain galactose, N-acetylgalactosamine and N-acetylneuraminic acid.

## Cellular localization

Secreted and Cell membrane.

## Images




Flow Cytometry - Anti-TNF alpha antibody [EPR20972] (Alexa Fluor® 647) (ab237354)

Overlay histogram showing RAW264.7 cells treated (red line) or untreated (green line) with 100 ng/mL of lipopolysaccharides (LPS) for 7 hours with addition of 1 µg/mL brefeldin A (BFA) for the last 3 hours and subsequent staining with ab237354. The cells were fixed with 4% formaldehyde and then permeabilized with 90% methanol. The cells were then incubated in 1x PBS containing 10% normal serum to block non-specific protein-protein interactions followed by the antibody (ab237354) ( $1 \times 10^6$  in 100µl at 1/500 dilution (1 µg/ml)) for 30 min at 22°C.

Isotype control antibody (black line) was Rabbit IgG (monoclonal) Alexa Fluor® 647 (ab199093) used at the same concentration and conditions. Unlabeled sample (blue line) was also used as a control.

Acquisition of >5,000 events were collected using a 17 mW Red laser (633 nm) and 660/20 bandpass filter.



Immunocytochemistry/ Immunofluorescence - Anti-TNF alpha antibody [EPR20972] (Alexa Fluor® 647) (ab237354)

ab237354 staining TNF alpha in RAW 264.7 +/- 100 ng/ml lipopolysaccharides (LPS) for 7 hours with addition of 1 µg/ml brefeldin A (BFA) for the last 3 hours. The cells were fixed with 4% formaldehyde (10 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab237354 at 1/100 dilution (shown in red) and ab195887, Mouse monoclonal to alpha Tubulin (Alexa Fluor® 488), at 1/250 dilution (shown in green). Nuclear DNA was labeled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

## **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors