

Product datasheet

Anti-Transcription factor AP-2-alpha antibody [EPR2688(2)] - BSA and Azide free ab236043

KO VALIDATED Recombinant RabMAB

10 Images

Overview

Product name	Anti-Transcription factor AP-2-alpha antibody [EPR2688(2)] - BSA and Azide free
Description	Rabbit monoclonal [EPR2688(2)] to Transcription factor AP-2-alpha - BSA and Azide free
Host species	Rabbit
Tested applications	Suitable for: Flow Cyt (Intra), IHC-P, ICC/IF, WB, IP
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
Positive control	IP: HeLa whole cell lysate; Flow Cyt (intra): JAR cells; ICC: JAR cells; IHC-P: Human breast carcinoma, and mouse and rat breast tissue; WB: HeLa, C6, Mouse skin and HAP1 cell lysates.
General notes	<p>ab236043 is the carrier-free version of ab108311.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAB[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAB[®] patents.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze.
Storage buffer	pH: 7.2 Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR2688(2)
Isotype	IgG

Applications

The Abpromise guarantee Our [Abpromise guarantee](#) covers the use of ab236043 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt (Intra)		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol.
ICC/IF		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration. Predicted molecular weight: 48 kDa.
IP		Use at an assay dependent concentration.

Target

Function Sequence-specific DNA-binding protein that interacts with inducible viral and cellular enhancer elements to regulate transcription of selected genes. AP-2 factors bind to the consensus sequence 5'-GCCNNNGGC-3' and activate genes involved in a large spectrum of important biological functions including proper eye, face, body wall, limb and neural tube development. They also suppress a number of genes including MCAM/MUC18, C/EBP alpha and MYC. AP-2-alpha is the only AP-2 protein required for early morphogenesis of the lens vesicle.

Involvement in disease Defects in TFAP2A are the cause of branchiooculofacial syndrome (BOFS) [MIM:113620]; also known as branchial clefts with characteristic facies, growth retardation, imperforate nasolacrimal duct, and premature aging or lip pseudocleft-hemangiomas branchial cyst syndrome. BOFS is a rare autosomal dominant cleft palate craniofacial disorder with variable expressivity. The major features include cutaneous anomalies, ocular anomalies, characteristic facial appearance (malformed pinnae, oral clefts), and, less commonly, renal and ectodermal (dental and hair)

anomalies.

Sequence similarities

Belongs to the AP-2 family.

Domain

The WW-binding motif mediates interaction with WWOX.


Post-translational modifications

Sumoylated on Lys-10; which inhibits transcriptional activity.

Cellular localization

Nucleus.

Images



All lanes : Anti-Transcription factor AP-2-alpha antibody [EPR2688(2)] ([ab108311](#)) at 1/10000 dilution (Purified)

Lane 1 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate

Lane 2 : C6 (Rat glial tumor glial cell) whole cell lysate

Lane 3 : Mouse skin lysate


Lysates/proteins at 15 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 48 kDa


This data was developed using [ab108311](#), the same antibody clone in a different buffer formulation.



Immunocytochemistry - Anti-Transcription factor AP-2-alpha antibody [EPR2688(2)] - BSA and Azide free (ab236043)

This data was developed using [ab108311](#), the same antibody clone in a different buffer formulation.


Immunocytochemistry analysis of JAR (Human placenta choriocarcinoma epithelial cell) cells labeling Transcription factor AP-2-alpha with Purified [ab108311](#) at 1/50 dilution (3.4 µg/mL). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/200 dilution (2.5 µg/mL). Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) was used as the secondary antibody at 1/1000 dilution (2 µg/mL). DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Transcription factor AP-2-alpha antibody [EPR2688(2)] - BSA and Azide free (ab236043)

This data was developed using [ab108311](#), the same antibody clone in a different buffer formulation.


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat breast tissue sections labeling Transcription factor AP-2-alpha with Purified [ab108311](#) at 1/100 dilution (1.07 µg/mL). Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0). Rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Transcription factor AP-2-alpha antibody [EPR2688(2)] - BSA and Azide free (ab236043)

This data was developed using [ab108311](#), the same antibody clone in a different buffer formulation.


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse breast tissue sections labeling Transcription factor AP-2-alpha with Purified [ab108311](#) at 1/100 dilution (1.07 µg/mL). Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0). Rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Transcription factor AP-2-alpha antibody [EPR2688(2)] - BSA and Azide free (ab236043)

This data was developed using [ab108311](#), the same antibody clone in a different buffer formulation.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human breast carcinoma tissue sections labeling Transcription factor AP-2-alpha with Purified [ab108311](#) at 1/100 dilution (1.07 µg/mL). Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0). Rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunoprecipitation - Anti-Transcription factor AP-2-alpha antibody [EPR2688(2)] - BSA and Azide free (ab236043)

This data was developed using [ab108311](#), the same antibody clone in a different buffer formulation.

Purified [ab108311](#) at 1/20 dilution (0.5µg) immunoprecipitating Transcription factor AP-2-alpha in HeLa whole cell lysate.

Lane 1 (input): HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate 10µg

Lane 2 (+): [ab108311](#) + HeLa whole cell lysate.


Lane 3 (-): Rabbit monoclonal IgG ([ab172730](#)) instead of [ab108311](#) in HeLa whole cell lysate.

VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) (1/10,000 dilution) was used for Western blotting.

Blocking Buffer and concentration: 5% NFDm/TBST.

Diluting buffer and concentration: 5% NFDm/TBST.


Observed band size: 48 kDa



Flow Cytometry (Intracellular) - Anti-Transcription factor AP-2-alpha antibody [EPR2688(2)] - BSA and Azide free (ab236043)

This data was developed using [ab108311](#), the same antibody clone in a different buffer formulation. Intracellular Flow Cytometry analysis of JAR (Human placenta choriocarcinoma epithelial cell) cells

labeling Transcription factor AP-2-alpha with Purified [ab108311](#) at 1/20 dilution (10 µg/mL) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Western blot - Anti-Transcription factor AP-2-alpha antibody [EPR2688(2)] - BSA and Azide free (ab236043)

All lanes : Anti-Transcription factor AP-2-alpha antibody [EPR2688(2)] ([ab108311](#)) at 1/1000 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : TFAP2A knockout HeLa cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.


Predicted band size: 48 kDa

Observed band size: 48 kDa

This data was developed using the same antibody clone in a different buffer formulation ([ab108311](#)).

Lanes 1-2: Merged signal (red and green). Green - [ab108311](#) observed at 48 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) observed at 37 kDa.

[ab108311](#) was shown to react with Transcription factor AP-2-alpha in wild-type HeLa cells in western blot. Loss of signal was observed when knockout cell line [ab265122](#) (knockout cell lysate [ab257736](#)) was used. Wild-type HeLa and TFAP2A knockout HeLa cell lysates were subjected to SDS-PAGE. [ab108311](#) and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) were incubated overnight at 4°C at a 1 in 1000 Dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-Transcription factor AP-2-alpha antibody [EPR2688(2)] - BSA and Azide free (ab236043)

All lanes : Anti-Transcription factor AP-2-alpha antibody [EPR2688(2)] ([ab108311](#)) at 1/1000 dilution

Lane 1 : Wild-type HAP1 whole cell lysate

Lane 2 : TFAP2A (AP2A) knockout HAP1 whole cell lysate

Lysates/proteins at 20 µg per lane.

Predicted band size: 48 kDa

Lanes 1 - 2: Merged signal (red and green). Green - [ab108311](#) observed at 48 kDa. Red - loading control, [ab9484](#), observed at 37 kDa.

[ab108311](#) was shown to recognize 0 in wild-type HAP1 cells as signal was lost at the expected MW in TFAP2A (AP2A) knockout cells. Additional cross-reactive bands were observed in the wild-type and knockout cells. Wild-type and TFAP2A (AP2A) knockout samples were subjected to SDS-PAGE. Ab108311 and [ab9484](#) (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed [ab216773](#) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed [ab216776](#) secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab108311](#)).

Why choose a recombinant antibody?

 Research with confidence Consistent and reproducible results	 Long-term and scalable supply Recombinant technology
 Success from the first experiment Confirmed specificity	 Ethical standards compliant Animal-free production

Anti-Transcription factor AP-2-alpha antibody [EPR2688(2)] - BSA and Azide free (ab236043)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors