

Product datasheet

Tyrosine Hydroxylase peptide ab183303

Description

Product name	Tyrosine Hydroxylase peptide
Animal free	No
Nature	Synthetic

Specifications

Our [Abpromise guarantee](#) covers the use of **ab183303** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	Blocking - Blocking peptide for Anti-Tyrosine Hydroxylase antibody [EP1533Y] (ab75875)
Form	Liquid
Additional notes	<ul style="list-style-type: none"> - First try to dissolve a small amount of peptide in either water or buffer. The more charged residues on a peptide, the more soluble it is in aqueous solutions. - If the peptide doesn't dissolve try an organic solvent e.g. DMSO, then dilute using water or buffer. - Consider that any solvent used must be compatible with your assay. If a peptide does not dissolve and you need to recover it, lyophilise to remove the solvent. - Gentle warming and sonication can effectively aid peptide solubilisation. If the solution is cloudy or has gelled the peptide may be in suspension rather than solubilised. - Peptides containing cysteine are easily oxidised, so should be prepared in solution just prior to use.

Preparation and Storage

Stability and Storage	Shipped at 4°C. Store at 4°C (stable for up to 12 months). Upon delivery aliquot. Store at -20°C long term.
------------------------------	---

General Info

Function	Plays an important role in the physiology of adrenergic neurons.
Tissue specificity	Mainly expressed in the brain and adrenal glands.
Pathway	Catecholamine biosynthesis; dopamine biosynthesis; dopamine from L-tyrosine: step 1/2.
Involvement in disease	Defects in TH are the cause of dystonia DOPA-responsive autosomal recessive (ARDRD)

[MIM:605407]; also known as autosomal recessive Segawa syndrome. ARDRD is a form of DOPA-responsive dystonia presenting in infancy or early childhood. Dystonia is defined by the presence of sustained involuntary muscle contractions, often leading to abnormal postures. Some cases of ARDRD present with parkinsonian symptoms in infancy. Unlike all other forms of dystonia, it is an eminently treatable condition, due to a favorable response to L-DOPA. Note=May play a role in the pathogenesis of Parkinson disease (PD). A genome-wide copy number variation analysis has identified a 34 kilobase deletion over the TH gene in a PD patient but not in any controls.

Sequence similarities

Belongs to the bipterin-dependent aromatic amino acid hydroxylase family.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors