

Product datasheet

Anti-XLF antibody [EPR15882-36] - C-terminal ab189917

KO VALIDATED Recombinant RabMAb[®]

1 References 7 Images

Overview

Product name	Anti-XLF antibody [EPR15882-36] - C-terminal
Description	Rabbit monoclonal [EPR15882-36] to XLF - C-terminal
Host species	Rabbit
Tested applications	Suitable for: Flow Cyt, ICC/IF, WB, IHC-P
Species reactivity	Reacts with: Human Does not react with: Mouse, Rat
Immunogen	Recombinant fragment within Human XLF aa 150 to the C-terminus. The exact sequence is proprietary. Database link: Q9H9Q4
Positive control	Ramos, Jurkat and HepG2 whole cell lysate (ab7900); Human endometrial adenocarcinoma; HepG2 and NCCIT cells; Ramos cells.

General notes

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information [see here](#).

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).

Reproducibility is key to advancing scientific discovery and accelerating scientists' next breakthrough.

Abcam is leading the way with our range of recombinant antibodies, knockout-validated antibodies and knockout cell lines, all of which support improved reproducibility.

We are also planning to innovate the way in which we present recommended applications and species on our product datasheets, so that only applications & species that have been tested in our own labs, our suppliers or by selected trusted collaborators are covered by our Abpromise[™] guarantee.

In preparation for this, we have started to update the applications & species that this product is

Abpromise guaranteed for.

We are also updating the applications & species that this product has been “predicted to work with,” however this information is not covered by our Abpromise guarantee.

Applications & species from publications and Abreviews that have not been tested in our own labs or in those of our suppliers are not covered by the Abpromise guarantee.

Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, as well as customer reviews and Q&As.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR15882-36
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab189917** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		1/60. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ICC/IF		1/250 - 1/500.
WB		1/10000 - 1/50000. Detects a band of approximately 39 kDa (predicted molecular weight: 33 kDa).
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target

Function	DNA repair protein involved in DNA nonhomologous end joining (NHEJ) required for double-strand break (DSB) repair and V(D)J recombination. May serve as a bridge between XRCC4 and the other NHEJ factors located at DNA ends, or may participate in reconfiguration of the end
-----------------	---

bound NHEJ factors to allow XRCC4 access to the DNA termini. It may act in concert with XRCC6/XRCC5 (Ku) to stimulate XRCC4-mediated joining of blunt ends and several types of mismatched ends that are noncomplementary or partially complementary.

Tissue specificity

Ubiquitously expressed.

Involvement in disease

Defects in NHEJ1 are the cause of severe combined immunodeficiency due to NHEJ1 deficiency (NHEJ1-SCID) [MIM:611291]; also known as autosomal recessive T cell-negative, B cell-negative, NK cell-positive, severe combined immunodeficiency with microcephaly, growth retardation and sensitivity to ionizing radiation or NHEJ1 syndrome. SCID refers to a genetically and clinically heterogeneous group of rare congenital disorders characterized by impairment of both humoral and cell-mediated immunity, leukopenia and low or absent antibody levels. Patients with SCID present in infancy with recurrent, persistent infections by opportunistic organisms. The common characteristic of all types of SCID is absence of T-cell-mediated cellular immunity due to a defect in T-cell development. NHEJ1-SCID is characterized by a profound T- and B-lymphocytopenia associated with increased cellular sensitivity to ionizing radiation, microcephaly and growth retardation. Some patients may manifest SCID with sensitivity to ionizing radiation without microcephaly and mild growth retardation, probably due to hypomorphic NHEJ1 mutations.

Note=A chromosomal aberration involving NHEJ1 is found in a patient with polymicrogyria. Translocation t(2;7)(q35;p22).

Sequence similarities

Belongs to the XLF family.

Cellular localization


Nucleus.

Images

Why choose a recombinant antibody?

- Research with confidence**
Consistent and reproducible results
- Long-term and scalable supply**
Recombinant technology
- Success from the first experiment**
Confirmed specificity
- Ethical standards compliant**
Animal-free production

Anti-XLF antibody [EPR15882-36] - C-terminal (ab189917)



Western blot - Anti-XLF antibody [EPR15882-36] - C-terminal (ab189917)


Lane 1: Wild type HAP1 whole cell lysate (20 µg)

Lane 2: XLF knockout HAP1 whole cell lysate (20 µg)

Lane 3: HeLa whole cell lysate (20 µg)


Lanes 1 - 3: Merged signal (red and green). Green - ab189917 observed at 38 kDa. Red - loading control, [ab18058](#), observed at 130 kDa.

ab189917 was shown to specifically react with XLF in wild type cells as signal was lost in XLF knockout cells. Wild-type and XLF knockout samples were subjected to SDS-PAGE. ab189917 and [ab18058](#) (Mouse anti-Vinculin loading control) were incubated overnight at 4°C at 1/10000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed [ab216773](#) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed [ab216776](#) secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.




Flow Cytometry - Anti-XLF antibody [EPR15882-36] - C-terminal (ab189917)

Flow Cytometrical analysis of Ramos cells labeling XLF with ab189917 at 1/60 compared to a negative control cell. FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Immunofluorescent analysis of paraformaldehyde-fixed NCCIT cells labeling XLF with ab189917 at 1/250, Goat anti rabbit IgG (Alexa Fluor® 555) at 1/200 and DAPI staining (blue).

Immunocytochemistry/ Immunofluorescence - Anti-XLF antibody [EPR15882-36] - C-terminal (ab189917)



Western blot - Anti-XLF antibody [EPR15882-36] - C-terminal (ab189917)

All lanes : Anti-XLF antibody [EPR15882-36] - C-terminal (ab189917) at 1/10000 dilution

Lane 1 : Ramos cell lysate

Lane 2 : Jurkat cell lysate

Lane 3 : HepG2 cell lysate


Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution


Predicted band size: 33 kDa

Additional bands at: 39 kDa. We are unsure as to the identity of these extra bands.



Immunofluorescent analysis of paraformaldehyde-fixed HepG2 cells labeling XLF with ab189917 at 1/250, Goat anti rabbit IgG (Alexa Fluor® 555) at 1/200 and DAPI staining (blue).

Immunocytochemistry/ Immunofluorescence - Anti-XLF antibody [EPR15882-36] - C-terminal (ab189917)



Immunohistochemical analysis of paraffin-embedded Human endometrial adenocarcinoma tissue labeling XLF with ab189917 at 1/250 with prediluted ImmunoHistoprobe(Ready to use) HRP Polymer for Rabbit IgG as secondary antibody.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-XLF antibody [EPR15882-36] - C-terminal (ab189917)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors